

Proceedings of EQUADIFF 2017 Conference



Editors:

Karol Mikula, Daniel Ševčovič, and Jozef Urbán

Published by Slovak University of Technology, 2017

SPEKTRUM
STU

This volume contains Proceedings of EQUADIFF 2017 Conference held in Bratislava, Slovakia, July 24-28, 2017, and organized by Slovak University of Technology in Bratislava, in cooperation with Comenius University in Bratislava, Union of Slovak Mathematicians and Physicists, Slovak Mathematical Society and Algoritmy:SK, Ltd.

© 2017, Published by Slovak University of Technology, SPEKTRUM STU Publishing

Karol Mikula, Daniel Ševčovič, and Jozef Urbán,
Editors of Proceedings of Equadiff 2017 Conference

ISBN 978-80-227-4757-8



Preface

The Equadiff is a series of biannual conferences on mathematical analysis, numerical approximation and applications of differential equations. It is held in rotation in the Czech Republic, Slovakia and Western Europe. The last Equadiff (Equadiff 14 in the Czecho-Slovak series) was organized in Bratislava, Slovakia, July 24-28, 2017 by the Slovak University of Technology, in cooperation with Comenius University, Union of Slovak Mathematicians and Physicists, Slovak Mathematical Society and Algoritmy:SK, ltd.

During the last decades the Equadiff has clearly developed into the world platform for international exchange of ideas on all mathematical and numerical aspects of differential equations, ranging from fundamental concepts to applications.

The scientific program of Equadiff 2017 Conference was proposed and prepared by the members International Scientific Programme Committee: Michal Beneš (Czech Technical University, Prague, Czech Republic), Charlie Elliott (University of Warwick, UK), Eduard Feireisl (Czech Academy of Sciences, Prague, Czech Republic), Marek Fila (Comenius University, Bratislava, Slovakia), Raphaele Herbin (University of Aix-Marseille, France), Grzegorz Karch (University of Wroclaw, Poland), Karol Mikula (Slovak University of Technology, Bratislava, Slovakia), Masayasu Mimura (Meiji University, Tokyo, Japan), Mario Ohlberger (University of Münster, Germany), Peter Poláčik (University of Minnesota, Minneapolis, USA), Otmar Scherzer (University of Vienna, Austria), Pavol Quittner (Comenius University, Bratislava, Slovakia), Eiji Yanagida (Tokyo Institute of Technology, Japan). Organizing Committee of the conference consisted of Peter Frolkovič, Angela Handlovičová, Martin Kalina, Karol Mikula, Daniel Ševčovič, Róbert Špir and Peter Struk. The conference was chaired by Karol Mikula and co-chaired by Marek Fila.

Proceedings of Equadiff 2017 Conference contain peer-reviewed contributions of participants of the conference. The proceedings cover a wide range of topics presented by plenary, minisymposia and contributed talks speakers. The scope of papers ranges from ordinary differential equations, differential inclusions and dynamical systems towards qualitative and numerical analysis of partial differential equations, stochastic PDEs and their applications.

In several papers, the authors studied qualitative and numerical properties of solutions to cross-diffusion systems with entropy structure, boundedness and stabilization of solutions in a three-dimensional and two-species chemotaxis-Navier-Stokes system, boundedness of solutions in a fully parabolic chemotaxis system with signal-dependent sensitivity and logistic term. Several authors studied well-posedness of solutions for a mass conserved Allen-Cahn equation with a nonlinear diffusion term, the porous medium equations and nonlinear cross-diffusion systems and efficient linear numerical scheme for solving the Stefan problem.

The authors also investigated qualitative behavior of solutions of the undamped Klein-Gordon equation and entropy of the attractor of the strongly damped wave equation. The conference proceedings contain papers on dynamical models of viscoplasticity and Lyapunov stability in hypoplasticity models. The proceedings further include papers dealing with qualitative properties of solutions for systems of fractional boundary value problems and analysis of inequalities with gradient nonlinearities and fractional Laplacian operators. The proceedings also contain papers dealing with qualitative properties like uniqueness and regularity of solutions for systems of coupled elliptic and parabolic equations.

Several papers are devoted to the numerical analysis of finite element and discrete Galerkin methods for elliptic problems with nonlinear boundary conditions. Applications of theoretical results cover viral infection modelling with diffusion and state-dependent delay, an analysis of a model of suspension flowing down an inclined plane as well as applications of tree-grid and finite stencil numerical methods in computational finance, optimal control and optimal design. Interesting applications of partial differential equations in image segmentation and computational differential geometry can be also found in the proceedings.

We thank all the authors for their interesting contributions to the conference proceedings. We also thank our reviewers for their valuable comments and suggestions which improved quality of presentation of results.

Bratislava, November 30, 2017

Karol Mikula, Daniel Ševčovič, and Jozef Urbán

Editors of Proceedings of Equadiff 2017 Conference

Table of Contents

Front matter	
Karol Mikula, Daniel Ševčovič, Jozef Urbán	i-viii
Positive solutions for a system of fractional boundary value problems	
Johnny Henderson, Rodica Luca	1-10
Boundedness and stabilization in a three-dimensional two-species chemotaxis-Navier-Stokes system	
Tomomi Yokota, Misaki Hirata, Shunsuke Kurima, Masaaki Mizukami	11-20
On the Solution Set of a Nonconvex Nonclosed Second Order Inclusion	
Aurelian Cernea	21-28
Dynamical model of viscoplasticity	
Konrad Kisiel	29-36
Nonlinear diffusion equations with perturbation terms on unbounded domains	
Shunsuke Kurima	37-44
On behavior of solutions to a chemotaxis system with a nonlinear sensitivity function	
Kentarou Fujie, Takasi Senba	45-52
Viral infection model with diffusion and state-dependent delay: a case of logistic growth	
Alexander V. Rezounenko	53-60
Boundedness in a fully parabolic chemotaxis system with signal-dependent sensitivity and logistic term	
Masaaki Mizukami	61-68
Kolmogorov's epsilon-entropy of the attractor of the strongly damped wave equation in locally uniform spaces	
Jakub Slavík	69-78
The Tree-Grid Method with Control-Independent Stencil	
Igor Kossaczky, Matthias Ehrhardt, Michael Günther	79-88

Multiple positive solutions for a p-Laplace critical problem ($p > 1$), via Morse theory	
Giuseppina Vannella	89-96
Propagation of errors in dynamic iterative schemes	
Barbara Zubik-Kowal	97-106
On Lyapunov stability in hypoplasticity	
Victor A. Kovtunenکو, Pavel Krejčí, Erich Bauer, Lenka Siváková, Anna V. Zubkova	107-116
Exponential convergence to the stationary measure and hyperbolicity of the minimisers for random Lagrangian	
Alexandre Boritchev	117-126
Analysis of the FEM and DGM for an elliptic problem with a nonlinear Newton boundary condition	
Miloslav Feistauer, Ondřej Bartoš, Filip Roskovec, Anna-Margarete Sändig	127-136
Numerical homogenization for indefinite H(curl)-problems	
Barbara Verfürth	137-146
Singularly perturbed set of periodic functional-differential equations arising in optimal control theory	
Valery Y. Glizer	147-156
Nonexistence of solutions of some inequalities with gradient nonlinearities and fractional Laplacian	
Evgeny Galakhov, Olga Salieva	157-162
Semi-analytical approach to initial problems for systems of nonlinear partial differential equations with constant delay	
Helena Šamajová	163-172
Visco-elastic-plastic modelling	
Jana Kopfová, Mária Minárová, Jozef Sumec	173-180
Cross-Diffusion Systems with Entropy Structure	
Ansgar Jüngel	181-190
Numerical modeling of heat exchange and unsaturated-saturated flow in porous media	
Jozef Kačur, Patrik Mihala, Michal Tóth	191-200
A well-posedness result for a mass conserved Allen-Cahn equation with nonlinear diffusion	
Perla El Kettani, Danielle Hilhorst, Kai Lee	201-210

Vectorial quasilinear diffusion equation with dynamic boundary condition	
Ryota Nakayashiki	211-220
Remarks on the qualitative behavior of the undamped Klein-Gordon equation	
Jorge A. Esquivel-Avila	221-228
Two approaches for the approximation of the nonlinear smoothing term in the image segmentation	
Matúš Tibenský, Angela Handlovičová	229-236
Stability of ALE space-time discontinuous Galerkin method	
Miloslav Vlasák, Monika Balázsová, Miloslav Feistauer	237-246
Upper Hausdorff dimension estimates for invariant sets of evolutionary systems on Hilbert manifolds	
Amina Kruck, Volker Reitman	247-254
Gaussian curvature based tangential redistribution of points on evolving surfaces	
Matej Medfa, Karol Mikula	255-264
Computational design optimization of low-energy buildings	
Jiří Vala	265-274
A generalization of the Keller-Segel system to higher dimensions from a structural viewpoint	
Kentarou Fujie, Takasi Senba	275-282
A Note on the Uniqueness and Structure of Solutions to the Dirichlet Problem for Some Elliptic Systems	
Jann-Long Chern, Shoji Yotsutani, Nichiro Kawano	283-286
Classical and generalized Jacobi polynomials orthogonal with different weight functions and differential equations satisfied by these polynomials	
Mariana Marčoková, Vladimír Guldan	287-294
Stochastic Modulation Equations on Unbounded Domains	
Luigi A. Bianchi, Dirk Blömker	295-304
An efficient linear numerical scheme for the Stefan problem, the porous medium equation and nonlinear cross-diffusion systems	
Motlatsi Molati, Hideki Murakawa	305-314
Continuous dependence for BV-entropy solutions to strongly degenerate parabolic equations with variable coefficients	
Hiroshi Watanabe	315-324

Numerical study on the blow-up rate to a quasilinear parabolic equation	
Koichi Anada, Tetsuya Ishiwata, Takeo Ushijima	325-330
Two methods for optical flow estimation	
Viera Kleinová, Peter Frolkovič	331-340
Mathematically Modelling The Dissolution Of Solid Dispersions	
Martin Meere, Sean McGint, Giuseppe Pontrelli	341-348
Toward a mathematical analysis for a model of suspension flowing down an inclined plane	
Kaname Matsue, Kyoko Tomoeda	349-358
Behaviour of the support of the solution appearing in some nonlinear diffusion equation with absorption	
Kenji Tomoeda	359-368
An elementary proof of asymptotic behavior of solutions	
Motohiro Sobajima, Giorgio Metafuno	369-376
Nonlinear Tensor Diffusion in Image Processing	
Oľga Stašová, Angela Handlovičová, Karol Mikula, Nadine Peyriéras	377-386
New efficient numerical method for 3D point cloud surface reconstruction by using level set methods	
Balázs Kósa, Jana Haličková-Brehovská, Karol Mikula	387-396
Converse problem for the two-component radial Gross-Pitaevskii system with a large coupling parameter	
Jean-Baptiste Casteras, Christos Sourdis	397-406

VIRAL INFECTION MODEL WITH DIFFUSION AND STATE-DEPENDENT DELAY: A CASE OF LOGISTIC GROWTH

ALEXANDER V. REZOUNENKO*

Abstract. We propose a virus dynamics model with reaction-diffusion and logistic growth terms, intracellular state-dependent delay and a general non-linear infection rate functional response. Classical solutions with Lipschitz in-time initial functions are investigated. This type of solutions is adequate to the discontinuous change of parameters due to, for example, drug administration. The Lyapunov functions approach is used to analyse stability of interior infection equilibria which describe the cases of a chronic disease.

Key words. Reaction-diffusion, evolution equations, Lyapunov stability, state-dependent delay, virus infection model.

AMS subject classifications. 93C23, 34K20,35K57, 97M60

1. Introduction. Our goal is to discuss a wide class of mathematical models of viral diseases. Many viruses (as Ebola virus, Zika virus, HIV, HBV, HCV and others) continue to be a major global public health issues, according to World Health Organization. Particularly, from The Global hepatitis report (WHO, April 2017) [25] we know that "a large number of people - about 325 million worldwide in 2015 - are carriers of hepatitis B or C virus infections, which can remain asymptomatic for decades." and "Viral hepatitis caused 1.34 million deaths in 2015, a number comparable to deaths caused by tuberculosis and higher than those caused by HIV. However, the number of deaths due to viral hepatitis is increasing over time, while mortality caused by tuberculosis and HIV is declining."

In such a situation any steps toward understanding the dynamics of viral diseases are important.

There are variety of models described by systems of ordinary differential equations and/or partial differential equations with or without delays which describe dynamics of different viral infections. Delays could be bounded or unbounded, concentrated or distributed, constant, time-dependent or state-dependent.

The classical models [12, 14] contain ordinary differential equations (without delay) for three variables: susceptible host cells T , infected host cells T^* and free virus particles V . The intracellular delay is an important property of the biological problem, so we start with the delay problem

$$(1.1) \quad \begin{cases} \dot{T}(t) = \lambda - dT(t) - f(T(t), V(t)), \\ \dot{T}^*(t) = e^{-\omega h} f(T(t-h), V(t-h)) - \delta T^*(t), \\ \dot{V}(t) = N\delta T^*(t) - cV(t). \end{cases}$$

In system (1.1), susceptible cells T are produced at a rate λ , die at rate dT , and become infected at rate $f(T, V)$. Properties and examples of incidence function f are discussed below. Infected cells T^* die at rate δT^* , free virions V are produced by infected cells at rate $N\delta T^*$ and are removed at rate $cV(t)$. In (1.1) h denotes the delay

*V.N.Karazin Kharkiv National University, Kharkiv, 61022, Ukraine (rezounenko@gmail.com) and Institute of Information Theory and Automation, Academy of Sciences of the Czech Republic, P.O. Box 18, 18208 Praha, CR

between the time a virus particle contacts a target cell and the time the cell becomes actively infected (start producing new *free* virions). It is clear that the constancy of the delay is just an extra assumption which essentially simplifies the study, but has no biological background.

To the best of our knowledge, viral infection models with state-dependent delay (SDD) have been considered for the first time in [20] (see also [21]). It is well known that differential equations with discrete state-dependent delay are always non-linear by its nature (see the review [5] for more details and discussion).

As usual in the study of delay systems with (maximal) delay $h > 0$ [4, 8], for a function $v(t), t \in [a - h, b] \subset \mathbf{R}, b > a$, we denote the history segment $v_t = v_t(\theta) \equiv v(t + \theta), \theta \in [-h, 0], t \in [a, b]$.

Consider a connected bounded domain $\Omega \subset \mathbf{R}^n$ with a smooth boundary $\partial\Omega$. Let $T(t, x), T^*(t, x), V(t, x)$ represent the densities of uninfected cells, infected cells and free virions at position $x \subset \Omega$ at time t .

In [22] the following system with SDD η is investigated

$$(1.2) \quad \begin{cases} \dot{T}(t, x) = \lambda - dT(t, x) - f(T(t, x), V(t, x)) + d^1 \Delta T(t, x), \\ \dot{T}^*(t, x) = e^{-\omega h} f(T(t - \eta(u_t), x), V(t - \eta(u_t), x)) - \delta T^*(t, x) + d^2 \Delta T^*(t, x), \\ \dot{V}(t, x) = N \delta T^*(t, x) - cV(t, x) + d^3 \Delta V(t, x). \end{cases}$$

Here the dot over a function denotes the partial time derivative i.g. $\dot{T}(t, x) = \frac{\partial T(t, x)}{\partial t}$, all the constants $\lambda, d, \delta, N, c, \omega$ are positive while $d^i, i = 1, 2, 3$ (diffusion coefficients) are non negative. In (1.2) (and in (1.3) below), a solution denoted by $u(t) = u(t, \cdot) = (T(t, \cdot), T^*(t, \cdot), V(t, \cdot))$, see the argument of the state-dependent delay η in the second equation. The precise definition of a solution is given below (Def. 2.1).

We consider a general functional response $f(T, V)$ satisfying natural assumptions presented below. In earlier models (with constant or without delay) the study was started in case of bilinear $f(T, V) = \text{const} \cdot TV$ and then extended to more general classes of non-linearities. For more details and discussion see [1, 3, 7, 11, 22].

We mention that the term $e^{-\omega h}$ in front of f (see the second equation (1.2)), in fact, states that only a *part* of the cell population survived during the virus incubation period. Clearly, it should be less than 1. It is an assumption which is not too precise in *nonlinear systems*. It could be regarded as a coefficient (strictly smaller than 1) and could be easily incorporated into the definition of the function f . We keep this coefficient in the form of $e^{-\omega h}$ for the only reason to simplify for the reader the comparison of computations with the constant delay case.

In this note we are interested in the following PDEs system with state-dependent delay η

$$(1.3) \quad \begin{cases} \dot{T}(t, x) = rT(t, x) \left(1 - \frac{T(t, x)}{T_K}\right) - dT(t, x) - f(T(t, x), V(t, x)) + d^1 \Delta T(t, x), \\ \dot{T}^*(t, x) = e^{-\omega h} f(T(t - \eta(u_t), x), V(t - \eta(u_t), x)) - \delta T^*(t, x) + d^2 \Delta T^*(t, x), \\ \dot{V}(t, x) = N \delta T^*(t, x) - cV(t, x) + d^3 \Delta V(t, x). \end{cases}$$

Let us discuss the principal difference in the first equations of (1.2) and (1.3). In system (1.2), uninfected target cells T are produced by the body at a constant rate λ which is relevant, for example, in case of HIV. In contrast, the first term in the first equation of (1.3) is the classical logistic growth term (Pierre Verhulst term) for the population of uninfected cells T . The constant T_K is the so-called carrying capacity for the population T , which has the clear biological meaning. System (1.3) is more

relevant in case of chronic infections with viruses such as, for example, hepatitis B (HBV) and hepatitis C (HCV). Here $T(t, x)$ and $T^*(t, x)$ represent uninfected and infected liver cells (hepatocytes). The carrying capacity could be also considered for the sum of uninfected and infected cells (c.f. [6]), but we decide to use it for uninfected hepatocytes (liver cells) only for the following biological reason. It is well-known that the development of HBV, HCV infections is usually connected with development of fibrosis. The last indicates that the regeneration of healthy hepatocytes is not quick enough to fill all the available (free) space in liver. This available space appears as a result of natural death of both uninfected and infected hepatocytes as well as killing of infected cells by immune system. The above suggests that the presence of infected cells does not make essential restriction on the regeneration of healthy hepatocytes T .

Boundary conditions are of Neumann type for the corresponding unknown if $d^i \neq 0$ i.e. $\frac{\partial T(t, x)}{\partial n}|_{\partial\Omega} = 0$ if $d^1 \neq 0$ and similarly for $T^*(t, x)$ and $V(t, x)$. Here $\frac{\partial}{\partial n}$ is the outward normal derivative on $\partial\Omega$. In case $d^i = 0$, no boundary conditions are needed for the corresponding unknown(s). For more discussion see [22].

Our main goals are to present the existence and uniqueness results for the model (1.3) in the sense of classical solutions, and to study the local asymptotic stability of non-trivial disease equilibria. We apply the Lyapunov approach [9] to the state-dependent delay PDE model (1.3) and allow, but not require, diffusion terms in each state equation. For the Lyapunov approach in context of viral infection models (with constant delay or nondelay cases) see e.g. works by A.Korobeinikov, C.McCluskey [7, 11] and references therein. Our main interest is in discussion of the state-dependent delay.

2. Main results. We use the basic functional framework described in [10] and applied to the system (1.2) in [22].

Define the following linear operator $-\mathcal{A}^0 = \text{diag}(d^1\Delta, d^2\Delta, d^3\Delta)$ in $C(\bar{\Omega}; \mathbf{R}^3)$ with $D(\mathcal{A}^0) \equiv D(d^1\Delta) \times D(d^2\Delta) \times D(d^3\Delta)$. Here, for $d^i \neq 0$ we set $D(d^i\Delta) \equiv \{v \in C^2(\bar{\Omega}) : \frac{\partial v(x)}{\partial n}|_{\partial\Omega} = 0\}$ and $D(d^j\Delta) \equiv C(\bar{\Omega})$ for $d^j = 0$. We omit the space coordinate x , for short, for unknown $u(t) = (T(t), T^*(t), V(t)) \in X \equiv [C(\bar{\Omega})]^3 \equiv C(\bar{\Omega}; \mathbf{R}^3)$. It is well-known that the closure $-\mathcal{A}$ (in X) of the operator $-\mathcal{A}^0$ generates a C_0 -semigroup $e^{-\mathcal{A}t}$ on X which is analytic and nonexpansive [10, p.5]. We denote the space of continuous functions by $C \equiv C([-h, 0]; X)$ equipped with the sup-norm $\|\psi\|_C \equiv \max_{\theta \in [-h, 0]} \|\psi(\theta)\|_X$.

We write, the system (1.3) in the following abstract form

$$(2.1) \quad \frac{d}{dt}u(t) + \mathcal{A}u(t) = F(u_t), \quad t > 0.$$

The non-linear continuous mapping $F : C \rightarrow X$ is defined by

$$(2.2) \quad F(\varphi)(x) = \begin{pmatrix} r\varphi^1(0, x) \left(1 - \frac{\varphi^1(0, x)}{T_K}\right) - d\varphi^1(0, x) - f(\varphi^1(0, x), \varphi^3(0, x)) \\ e^{-\omega h} f(\varphi^1(-\eta(\varphi), x), \varphi^3(-\eta(\varphi), x)) - \delta\varphi^2(0, x) \\ N\delta\varphi^2(0, x) - c\varphi^3(0, x) \end{pmatrix}.$$

Here $\varphi = (\varphi^1, \varphi^2, \varphi^3) \in C$. Mapping F is *not* Lipschitz on the space C which is typical for a mapping which includes discrete state-dependent delays (see review [5] for ODE case and works [15, 16, 17, 2] for PDEs).

We need initial conditions $u(\theta, x) = \varphi(\theta, x) = (T(\theta, x), T^*(\theta, x), V(\theta, x)), \theta \in [-h, 0]$ for the delay problem (2.1) (c.f. (1.3)):

$$(2.3) \quad \varphi \in Lip([-h, 0]; X) \equiv \left\{ \psi \in C : \sup_{s \neq t} \frac{\|\psi(s) - \psi(t)\|_X}{|s - t|} < \infty \right\}, \quad \varphi(0) \in D(A).$$

In our study we use the standard (c.f. [13, Def. 2.3, p.106] and [13, Def. 2.1, p.105])

DEFINITION 2.1. *A function $u \in C([-h, T]; X)$ is called a **mild solution** on $[-h, T)$ of the initial value problem (2.1), (2.3) if it satisfies (2.3) and $u(t) = e^{-At}\varphi(0) + \int_0^t e^{-A(t-s)}F(u_s) ds$, $t \in [0, T)$.*

*A function $u \in C([-h, T]; X) \cap C^1((0, T); X)$ is called a **classical solution** on $[-h, T)$ of the initial value problem (2.1), (2.3) if it satisfies (2.3), $u(t) \in D(A)$ for $0 < t < T$ and (2.1) is satisfied on $(0, T)$.*

Assume the non-linear function $f : \mathbf{R}^2 \rightarrow \mathbf{R}$ is Lipschitz continuous and satisfies

$$(2.4) \quad (\mathbf{Hf}_1) \quad \text{there exists } \mu > 0 \text{ such that } |f(T, V)| \leq \mu|T| \text{ for all } T, V \in \mathbf{R}.$$

We have the following result

THEOREM 2.2. *Let nonlinear function f be Lipschitz and satisfy (\mathbf{Hf}_1) (see (2.4)), state-dependent delay $\eta : C \rightarrow [0, h]$ is locally Lipschitz. Then the initial value problem (2.1), (2.3) has a unique classical solution which is global in time i.e. defined for all $t \geq 0$.*

Proof of Theorem 2.2 follows the line of the proof of [22, Proposition 1].

Define the set (c.f. (2.3)), which is different from the one Ω_{Lip} in [22]:

$$\begin{aligned} \Omega_{Lip}^{log} &\equiv \{ \varphi = (\varphi^1, \varphi^2, \varphi^3) \in Lip([-h, 0]; X) \subset C, \varphi(0) \in D(A) : \\ &0 \leq \varphi^1(\theta) \leq \left(1 - \frac{d}{r}\right) T_K, 0 \leq \varphi^2(\theta) \leq \frac{\mu}{\delta} \left(1 - \frac{d}{r}\right) T_K e^{-\omega h}, \\ (2.5) \quad &0 \leq \varphi^3(\theta) \leq \frac{N\mu}{c} \left(1 - \frac{d}{r}\right) T_K e^{-\omega h}, \quad \theta \in [-h, 0] \}, \end{aligned}$$

where μ is defined in (\mathbf{Hf}_1) and all the inequalities hold pointwise w.r.t. $x \in \bar{\Omega}$.

We need further assumptions (which include (\mathbf{Hf}_1)) on Lipschitz function f :

$$(2.6) \quad (\mathbf{Hf}_1+) \quad \begin{cases} f(T, 0) = f(0, V) = 0, \quad \text{and} \quad f(T, V) > 0 \text{ for all } T > 0, V > 0; \\ f \text{ is strictly increasing in both coordinates for all } T > 0, V > 0; \\ \text{there exists } \mu > 0 \text{ such that } |f(T, V)| \leq \mu|T| \text{ for all } T, V \in \mathbf{R}. \end{cases}$$

We have the following result

THEOREM 2.3. *Let non-linear Lipschitz function f satisfy (\mathbf{Hf}_1+) (see (2.6)), state-dependent delay $\eta : C \rightarrow [0, h]$ is locally Lipschitz. Then Ω_{Lip}^{log} is invariant i.e. for any $\varphi \in \Omega_{Lip}^{log}$ the unique solution to problem (2.1), (2.3) satisfies $u_t \in \Omega_{Lip}^{log}$ for all $t \geq 0$.*

Proof of Theorem 2.3. The existence and uniqueness of solution is proven in theorem 2.2. The proof of the invariance part follows the invariance result of [10] with the use of the *almost Lipschitz property* of nonlinearity F . The estimates (for the subtangential condition) are the same as for the constant delay case, see e.g. [11, Theorem 2.2]. We do not repeat it here. It is important to notice that the solutions are classic for all $t \geq 0$ (but not for $t \geq h$ as could be in the case of merely continuous initial functions $\varphi \in C$). For more details see, e.g. [22]. The proof of Theorem 2.3 is complete.

2.1. Stationary solutions. Let us discuss stationary solutions of (1.3). By such solutions we mean time independent \widehat{u} which, in general, may depend on $x \in \overline{\Omega}$. Consider the system (1.3) with $u(t) = u(t - \eta(u_t)) = \widehat{u}$ and denote the coordinates (a possible triple of coordinates) of a stationary solution by $(\widehat{T}, \widehat{T}^*, \widehat{V}) = \widehat{u} \equiv \widehat{\varphi}(\theta)$, $\theta \in [-h, 0]$. Since stationary solutions of (1.3) do not depend on the type of delay (state-dependent or constant) we have

$$(2.7) \quad \begin{cases} 0 = r\widehat{T} \left(1 - \frac{\widehat{T}}{T_K}\right) - d\widehat{T} - f(\widehat{T}, \widehat{V}), & 0 = e^{-\omega h} f(\widehat{T}, \widehat{V}) - \delta\widehat{T}^*, \\ 0 = N\delta\widehat{T}^* - c\widehat{V}. \end{cases}$$

Equations hold pointwise w.r.t. $x \in \overline{\Omega}$.

It is easy to see that the trivial stationary solution $((1 - \frac{d}{r})T_K, 0, 0)$ always exists. We are interested in nontrivial disease stationary solutions of (1.3). We have from the first and second equations of (2.7) $\widehat{T}^* = \frac{r}{\delta}e^{-\omega h} \cdot \widehat{T} \left(1 - \frac{\widehat{T}}{T_K}\right) - \frac{d}{\delta}e^{-\omega h}\widehat{T}$ and from the third equation $\widehat{V} = \frac{N\delta}{c}\widehat{T}^*$. It gives the condition on the coordinate \widehat{T} which should belong to $(0, (1 - \frac{d}{r})T_K]$. Denote (c.f. [11, 20])

$$(2.8) \quad \begin{aligned} h_f^{log}(s) \equiv & f\left(s, \frac{Nr}{c}e^{-\omega h} \cdot s \left(1 - \frac{s}{T_K}\right) - \frac{Nd}{c}e^{-\omega h} \cdot s\right) \\ & - r \cdot s \left(1 - \frac{s}{T_K}\right) + d \cdot s. \end{aligned}$$

Assume f satisfies

$$(\mathbf{Hf}_2^{\log}) \quad h_f^{log}(s) = 0 \text{ has at least one and at most a finite number} \\ \text{of roots on } (0, (1 - \frac{d}{r})T_K].$$

We denote an arbitrary root of $h_f^{log}(s) = 0$ by \widehat{T} and define the corresponding $\widehat{T}^* \equiv \frac{r}{\delta}e^{-\omega h} \cdot \widehat{T} \left(1 - \frac{\widehat{T}}{T_K}\right) - \frac{d}{\delta}e^{-\omega h}\widehat{T}$ and $\widehat{V} \equiv \frac{N\delta}{c}\widehat{T}^* = \frac{Nr}{c}e^{-\omega h} \cdot \widehat{T} \left(1 - \frac{\widehat{T}}{T_K}\right) - \frac{Nd}{c}e^{-\omega h}\widehat{T}$. The point $(\widehat{T}, \widehat{T}^*, \widehat{V})$ satisfies (2.7), so it is a disease stationary solution of (1.3). We notice that in [22] the corresponding equation was written for coordinate \widehat{T}^* , while (2.8) is designed for $s = \widehat{T}$.

Remark (c.f. [22]). *We notice that the finiteness of roots (which are obviously isolated) does not allow the existence of equilibria which depend on spatial coordinate $x \in \Omega$. We remind that Ω is a connected set, so a function $v \in C(\overline{\Omega})$ may take either one or continuum values. Assumption (\mathbf{Hf}_2^{\log}) implies $\widehat{T}^*(x) \equiv \widehat{T}^* \in \mathbf{R}$, so $(\widehat{T}, \widehat{T}^*, \widehat{V})$ is independent of $x \in \overline{\Omega}$.*

Remark. *It is important to mention that usually in study of stability properties of stationary solutions (for viral dynamics problems) one uses conditions on the so-called reproduction numbers. These conditions are used to separate the case of a unique stationary solution. Then the global stability of the equilibrium is investigated. In our study, taking into account the state-dependence of the delay, we discuss the local stability. As a consequence, it allows the co-existence of multiple equilibria. We believe this framework provides a way to model more complicated situations with rich dynamics (in contrast to a globally stable equilibrium). The conditions on the reproduction numbers do not appear explicitly here, but could be seen as particular sufficient conditions for (\mathbf{Hf}_2^{\log}) .*

2.2. Stability of disease stationary solutions. In this section we use the following *local* assumptions on f in a small neighborhood of a disease equilibrium (given by (\mathbf{Hf}_2^{\log})).

$$(2.9) \quad (\mathbf{Hf}_3) \quad \left(\frac{V}{\widehat{V}} - \frac{f(T, V)}{f(T, \widehat{V})} \right) \cdot \left(\frac{f(T, V)}{f(T, \widehat{V})} - 1 \right) > 0.$$

One can check that the DeAngelis-Beddington functional response [1, 3] of the form $f(T, V) = \frac{kTV}{1+k_1T+k_2V}$, with $k, k_1 \geq 0, k_2 > 0$ satisfies (\mathbf{Hf}_3) globally. We also mention that the DeAngelis-Beddington functional response includes as a special case ($k_1 = 0$) the *saturated incidence* rate $f(T, V) = \frac{kTV}{1+k_2V}$.

We will also use the assumption

(\mathbf{Hf}_4) Function f is differentiable in a neighborhood of $(\widehat{T}, \widehat{V})$.

The main result is the following

THEOREM 2.4. *Let the nonlinear Lipschitz function f satisfy (\mathbf{Hf}_1+) , (\mathbf{Hf}_2^{\log}) , (\mathbf{Hf}_3) , (\mathbf{Hf}_4) (see (2.6), (2.9)), a root \widehat{T} of $h_f^{\log}(s) = 0$ (see (2.8) and (\mathbf{Hf}_2^{\log})) satisfy $\widehat{T} > \frac{1}{2}(1 - \frac{d}{r})T_K$. Let state-dependent delay $\eta : C \rightarrow [0, h]$ be locally Lipschitz in C and continuously differentiable in a neighbourhood of equilibrium $\widehat{\varphi} \equiv (\widehat{T}, \widehat{T}^*, \widehat{V})$. Then the stationary solution $\widehat{\varphi}$ is locally asymptotically stable.*

In the proof we use the following Lyapunov functional with *state-dependent delay* along a solution of (1.3)

$$(2.10) \quad U^{\text{sdd}}(t) \equiv \int_{\Omega} \left\{ \left(T(t, x) - \widehat{T} - \int_{\widehat{T}}^{T(t, x)} \frac{f(\widehat{T}, \widehat{V})}{f(\theta, \widehat{V})} d\theta \right) e^{-\omega h} + \widehat{T}^* \cdot v \left(\frac{T^*(t, x)}{\widehat{T}^*} \right) + \frac{\widehat{V}}{N} \cdot v \left(\frac{V(t, x)}{\widehat{V}} \right) + \delta \widehat{T}^* \int_{t-\eta(u_t)}^t v \left(\frac{f(T(\theta, x), V(\theta, x))}{f(\widehat{T}, \widehat{V})} \right) d\theta \right\} dx.$$

In (2.10) the Volterra function $v(s) = s - 1 - \ln s : (0, +\infty) \rightarrow \mathbf{R}_+$ (c.f. [7, 11]) is used. The form of the functional is standard except the low limit of the last integral in (2.10) which is state-dependent. This state-dependence was first considered in [20] (see also [21]). For PDE (1.2) with constant delay case and $d^1 = d^2 = 0$, see e.g. [11] and for PDE with state-dependent delay (1.2) see [22]. We do not repeat here detailed calculations of the time derivative of $U^{\text{sdd}}(t)$ along a solution of (1.3). They are similar to the ones of [22] and differ in the parts where the connection between coordinates of the stationary solution $\widehat{\varphi} = (\widehat{T}, \widehat{T}^*, \widehat{V})$ is used. The logistic growth term also makes difference to the study presented in [20, 22].

Acknowledgments. The author is thankful to A.Korobeinikov for useful discussions. This work was supported in part by GA CR under project 16-06678S.

REFERENCES

- [1] J. R. BEDDINGTON, *Mutual interference between parasites or predators and its effect on searching efficiency*, *Journal of Animal Ecology*, **44** (1975), 331-340. pp.
- [2] I.D. CHUESHOV AND A.V. REZOUNENKO, *Finite-dimensional global attractors for parabolic nonlinear equations with state-dependent delay*, *Communications on Pure and Applied Analysis*, **14/5** (2015), pp.1685-1704.
- [3] D. L. DEANGELIS, R. A. GOLDSTEIN AND R. V. O'NEILL, *A model for tropic interaction*, *Ecology*, **56** (1975), pp.881-892.
- [4] J. K. HALE, *Theory of Functional Differential Equations*, Springer, Berlin- Heidelberg- New York, 1977.
- [5] F. HARTUNG, T. KRISZTIN, H.-O. WALTHER AND J. WU, *Functional differential equations with state-dependent delays: Theory and applications*, In: *Canada, A., Drabek., P. and A. Fonda (Eds.) Handbook of Differential Equations, Ordinary Differential Equations*, Elsevier Science B.V., North Holland, **3** (2006), pp.435-545.
- [6] S. HEWS, S. EIKENBERRY, J.D. NAGY ET AL., *Rich dynamics of a hepatitis B viral infection model with logistic hepatocyte growth*, *Journal of Mathematical Biology*, Volume 60, Issue 4, (2010), pp.573-590.
- [7] A. KOROBENIKOV, *Global properties of infectious disease models with nonlinear incidence*, *Bull. Math. Biol.*, **69** (2007), pp.1871-1886.
- [8] Y. KUANG, *Delay Differential Equations with Applications in Population Dynamics*, *Mathematics in Science and Engineering*, 191. Academic Press, Inc., Boston, MA, 1993.
- [9] A. M. LYAPUNOV, *The General Problem of the Stability of Motion*, Kharkov Mathematical Society, Kharkov, 1892, 251p.
- [10] R.H. MARTIN, JR., H.L. SMITH, *Abstract functional-differential equations and reaction-diffusion systems*, *Trans. Amer. Math. Soc.*, **321** (1990), pp.1-44.
- [11] C. MCCLUSKEY, YU.YANG, *Global stability of a diffusive virus dynamics model with general incidence function and time delay*, *Nonlinear Anal. Real World Appl*, **25** (2015), pp.64-78.
- [12] M. NOWAK AND C. BANGHAM, *Population dynamics of immune response to persistent viruses*, *Science*, **272** (1996), pp.74-79.
- [13] A. PAZY, *Semigroups of linear operators and applications to partial differential equations*, *Applied Mathematical Sciences*, **44**. Springer-Verlag, New York, 1983. viii+279 pp.
- [14] A. PERELSON, A. NEUMANN, M. MARKOWITZ, J. LEONARD AND D. HO, *HIV-1 dynamics in vivo: Virion clearance rate, infected cell life-span, and viral generation time*, *Science*, **271** (1996), pp.1582-1586.
- [15] A. V. REZOUNENKO, *Partial differential equations with discrete and distributed state-dependent delays*, *Journal of Mathematical Analysis and Applications*, **326** (2007), pp.1031-1045.
- [16] A. V. REZOUNENKO, *Differential equations with discrete state-dependent delay: Uniqueness and well-posedness in the space of continuous functions*, *Nonlinear Analysis: Theory, Methods and Applications*, **70** (2009), pp.3978-3986.
- [17] A. V. REZOUNENKO, *Non-linear partial differential equations with discrete state-dependent delays in a metric space*, *Nonlinear Analysis: Theory, Methods and Applications*, **73** (2010), pp.1707-1714.
- [18] A. V. REZOUNENKO, *A condition on delay for differential equations with discrete state-dependent delay*, *Journal of Mathematical Analysis and Applications*, **385** (2012), pp.506-516.
- [19] A.V. REZOUNENKO, P. ZAGALAK, *Non-local PDEs with discrete state-dependent delays: well-posedness in a metric space*, *Discrete and Continuous Dynamical Systems - Series A*, **33:2** (2013), pp.819-835.
- [20] A. V. REZOUNENKO, *Stability of a viral infection model with state-dependent delay, CTL and antibody immune responses*, *Discrete and Continuous Dynamical Systems - Series B*, Vol. **22** (2017), pp.1547-1563; Preprint arXiv:1603.06281v1 [math.DS], 20 March 2016, arxiv.org/abs/1603.06281v1.
- [21] A. V. REZOUNENKO, *Continuous solutions to a viral infection model with general incidence rate, discrete state-dependent delay, CTL and antibody immune responses*, *Electron. J. Qual. Theory Differ. Equ.*, **79** (2016), pp.1-15.
- [22] A. V. REZOUNENKO, *Viral infection model with diffusion and state-dependent delay: stability of classical solutions*, *Discrete and Continuous Dynamical Systems - Series B*, Vol. **23**, N. 3, May 2018, to appear; Preprint arXiv:1706.08620 [math.DS], 26 Jun 2017, arxiv.org/abs/1706.08620.
- [23] H. L. SMITH, *Monotone Dynamical Systems. An Introduction to the Theory of Competitive and Cooperative Systems*, *Mathematical Surveys and Monographs*, **41**. American Mathematical

- Society, Providence, RI, 1995.
- [24] H. SMITH, *An Introduction to Delay Differential Equations with Sciences Applications to the Life*, Texts in Applied Mathematics, vol. 57, Springer, New York, Dordrecht, Heidelberg, London, 2011.
- [25] WORLD HEALTH ORGANIZATION, *Global hepatitis report-2017, April 2017*, ISBN: 978-92-4-156545-5; apps.who.int/iris/bitstream/10665/255016/1/9789241565455-eng.pdf?ua=1