

GENERALIZED $W^{1,1}$ -YOUNG MEASURES AND RELAXATION OF PROBLEMS WITH LINEAR GROWTH*

MARGARIDA BAÍA[†], STEFAN KRÖMER[‡], AND MARTIN KRUŽÍK[§]

Abstract. In this work we completely characterize generalized Young measures generated by sequences of gradients of maps in $W^{1,1}(\Omega; \mathbb{R}^M)$, where $\Omega \subset \mathbb{R}^N$. This characterization extends and completes previous analysis by Kristensen and Rindler [*Arch. Ration. Mech. Anal.*, 197 (2010), pp. 539–598 and 203 (2012), pp. 693–700] where concentrations of the sequence of gradients at the boundary of Ω were excluded. As an application of our result we study the relaxation of non-quasiconvex variational problems with linear growth at infinity, and, finally, we link our characterization to Souček spaces [J. Souček, *Časopis Pro Pěstování Matematiky*, 97 (1972), pp. 10–46], an extension of $W^{1,1}(\Omega; \mathbb{R}^M)$ where gradients are considered as measures on $\bar{\Omega}$.

Key words. lower semicontinuity, BV, quasiconvexity, free boundary, Young measures and their generalizations

AMS subject classifications. 49J45, 26B30, 52A99

DOI. 10.1137/16M1103464

1. Introduction. Oscillations and concentrations appear in many problems of the calculus of variations, partial differential equations, and/or optimal control theory; cf. [6] for an overview and examples from elasticity and continuum mechanics of solids. While Young measures [35] successfully capture oscillatory behavior of sequences, they completely miss concentration effects. These concentration effects may be dealt with by appropriate generalizations of Young measures, as in DiPerna’s and Majda’s treatment of concentrations [10], following Alibert’s and Bouchitté’s approach in [1] (see also [11]), etc. Detailed overviews of this subject may be found in [29, 32]. A general feature of all these approaches is the description of oscillations and/or concentrations in sequences in terms of parametrized measures. Besides the work of Kinderlehrer and Pedregal [16, 17] dealing with Young measures, and thus with oscillations only, both concentrations and oscillations were jointly treated in [22] in $W^{1,p}$ for $p > 1$ and in [20, 28] in $W^{1,1}$ and BV. Applications of parametrized measures in optimal control of dynamical systems with linear growth involving oscillations and concentrations can be found, e.g., in [8, 25, 29].

Explicit characterization of such parametrized measures is very important for applications because it allows us to analyze limit problems. The parametrized measures generated by sequences of gradients are of a particular interest in vectorial multidimensional calculus of variations, for instance when we want to minimize integral

*Received by the editors November 14, 2016; accepted for publication (in revised form) September 26, 2017; published electronically February 13, 2018.

<http://www.siam.org/journals/sima/50-1/M110346.html>

Funding: The work of the authors was supported by GAČR through projects 14-15264S and 16-34894L. The first author acknowledges the Czech Academy of Sciences, Institute of Information Theory and Automation for its hospitality and support, and the third author is indebted to CAMGSD for the support by the project UID/MAT/04459/2013 during his stay there.

[†]CAMGSD/Departamento de Matemática, Instituto Superior Técnico, 1049-001 Lisbon, Portugal (mbaia@math.tecnico.ulisboa.pt).

[‡]The Czech Academy of Sciences, Institute of Information Theory and Automation, CZ-182 08 Praha 8, Czech Republic (skroemer@utia.cas.cz).

[§]The Czech Academy of Sciences, Institute of Information Theory and Automation, CZ-182 08 Praha 8, Czech Republic, and Faculty of Civil Engineering, Czech Technical University, CZ-166 29 Praha 6, Czech Republic (kruzik@utia.cas.cz).

functionals $I(u) := \int_{\Omega} v(\nabla u(x)) \, dx$ with a nonconvex (or non-quasiconvex) integrand v having p -growth at infinity, i.e., for large arguments. The functional F is then not lower semicontinuous with respect to weak convergence in $W^{1,p}(\Omega; \mathbb{R}^m)$ and its minimizers do not necessarily exist. Parametrized measures are then applied to define the so-called relaxed problem tracing the limit behavior of minimizing sequences. For sequences of gradients of maps living in $W^{1,p}(\Omega; \mathbb{R}^M)$, Kinderlehrer and Pedregal characterized Young measures in [16, 17] for $p = +\infty$ and $1 < p < +\infty$, respectively. If $p > 1$, concentrations effects in minimizing sequence of coercive functionals are excluded using the so-called decomposition lemma proved in [13] which allows one to construct minimizing sequences which do not exhibit concentrations. Consequently, Young measures suffice to describe the relaxed functional. On the other hand, concentration effects can appear if v is noncoercive, more precisely if $|v(A)| \leq C(1 + |A|^p)$ for some $C > 0$. Generalizations of Young measures were used in [14, 15, 23] to analyze necessary and sufficient conditions guaranteeing weak lower semicontinuity of F ; cf. also [6].

If $p = 1$, i.e, for functionals with linear growth, concentration effects are usually always relevant and often prevent the existence of minimizers in $W^{1,1}$, a well-known fact intimately related to the lack of reflexivity of $W^{1,1}$. For precisely that reason, the standard approach to handle functionals with linear growth relies on extension to BV, which in particular allows to keep track of certain concentrations of the gradient in the interior of the domain, as singular contributions of the derivative in BV. Moreover, if the integrand lacks (quasi)convexity, oscillations and concentrations appear simultaneously and are mutually interconnected. The methods used to characterize Young measures and their generalizations for $p > 1$ cannot then be applied; in particular, the decomposition lemma [13, Lemma 1.2], which allows us to replace a sequence of gradients generating a Young measure by a new sequence of gradients such that the p th power of the norm is equiintegrable, does not hold in $W^{1,1}$ or BV.

In [20], the authors resolved this problem *assuming* that concentrations do not appear at the boundary of the domain Ω . As also shown in [20], in particular this is enough to discuss weak* lower semicontinuity for integral functionals in BV subject to a Dirichlet condition, because it is then possible to extend both the integrand and the admissible functions to a larger domain $\tilde{\Omega}$, where everything will be fixed near the boundary. This of course excludes concentration effects near the boundary of $\partial\Omega$. On the other hand, as soon as there is at least a part of the boundary where we do not impose a Dirichlet condition, one cannot fully avoid this phenomenon as the following simple example illustrates.

Consider the following one-dimensional variational problem:

$$(1.1) \quad \min_{u \in W^{1,1}(0,1)} \left\{ I(u) := \int_0^1 ((x-1)^2 + \varepsilon)|u'(x)| \, dx + u(0)^2 + (u(1)-1)^2 \right\}$$

for given $0 < \varepsilon < 1$. Notice that although no Dirichlet boundary condition was imposed, we still get the following natural boundary conditions for critical points, at least formally:

$$(1 + \varepsilon) \frac{u'}{|u'|} \varrho + 2u = 0 \text{ at } x = 0 \quad \text{and} \quad \varepsilon \frac{u'}{|u'|} \varrho + 2(u - 1) = 0 \text{ at } x = 1$$

(nonlinear Robin condition),

where ϱ is the outer normal at the boundary: $\varrho(0) = -1$, $\varrho(1) = +1$.

Roughly speaking, members of a minimizing sequence $(u_n)_{n \in \mathbb{N}}$ for (1.1) will try to keep their derivative as small as possible but also need to achieve some values close

to zero and one at $x = 0$ and $x = 1$, respectively. Due to the weight $x \mapsto (x - 1)^2 + \varepsilon$, the best place to switch from one constant to another one is near $x = 1$. In fact, the infimum of I can be calculated explicitly, and we can also give an explicit minimizing sequence: For every u admissible,

$$I(u) > \varepsilon \int_0^1 |u'(x)| dx + u(0)^2 + (u(1) - 1)^2 \geq \varepsilon(u(1) - u(0)) + u(0)^2 + (u(1) - 1)^2 .$$

The function $(u(0), u(1)) \mapsto \varepsilon(u(1) - u(0)) + u(0)^2 + (u(1) - 1)^2$ is minimized for $u(0) = \varepsilon/2$ and $u(1) = (2 - \varepsilon)/2$ with the value $(2\varepsilon - \varepsilon^2)/2$. On the other hand, take

$$u_n(x) := \begin{cases} \varepsilon/2 & \text{if } 0 \leq x \leq 1 - 1/n, \\ n(1 - \varepsilon)x + \varepsilon/2 - (1 - \varepsilon)(n - 1) & \text{otherwise.} \end{cases}$$

We see that $I(u_n) \rightarrow (2\varepsilon - \varepsilon^2)/2$ as $n \rightarrow \infty$, so it is a minimizing sequence. However, $u_n \rightarrow 0$ in $L^1(0, 1)$ and (u'_n) concentrates at $x = 1$. In particular, no minimizer exists in the admissible class of competitors in $W^{1,1}(0, 1)$.

It is not entirely obvious how to relax I in BV, or other functionals (formally) involving a natural boundary condition of Robin type. In fact, if we simply replace the integral part of I with its relaxation with respect to weak* convergence in BV while leaving the boundary term alone, we get for all $u \in \text{BV}(0, 1)$

$$I_1(u) := \int_0^1 ((x - 1)^2 + \varepsilon) d|Du(x)| + u(0)^2 + (u(1) - 1)^2,$$

where $u(0)$ and $u(1)$ are understood in the sense of trace in BV. However, this clearly is not the right choice, because I_1 is still not lower semicontinuous with respect to weak* convergence in BV for $\varepsilon < 2$: with the minimizing sequence (u_n) defined above, we have $u_n \rightharpoonup^* 0$ in BV and $I_1(u_n) = I(u_n) \rightarrow (4\varepsilon - \varepsilon^2)/4 < I_1(0) = 1$. Now, the main issue is that the trace at $x = 1$ is not continuous along (u_n) , which here causes the term $(u_n(1) - 1)^2$ to jump up in the limit, reflecting the fact that the concentration of (u'_n) at $x = 1$ is being dropped in the weak* limit in BV. The natural way to solve this dilemma is to consider a slightly larger space, and a good choice is $X := \text{BV}(0, 1) \times \mathbb{R}^2$, the closure of $W^{1,1}$ (and BV), embedded into X by $u \in W^{1,1} \mapsto (u, (u(0), u(1))) \in X$, with respect to weak* convergence in X . Essentially, we now keep track of the “true” boundary value at $x = 0, 1$ along sequences in $W^{1,1}$ possibly different from the trace of the limit function in BV, and it is not difficult to check that the relaxation of I (and I_1) in X is given for all $(u, (\beta_0, \beta_1)) \in \text{BV}(0, 1) \times \mathbb{R}^2$ by

$$I_2(u, (\beta_0, \beta_1)) := \int_0^1 ((x - 1)^2 + \varepsilon) d|Du(x)| + (1 + \varepsilon)|u(0) - \beta_0| + \beta_0^2 + \varepsilon|\beta_1 - u(1)| + (\beta_1 - 1)^2,$$

where $u(0)$ and $u(1)$ are understood in the sense of traces in BV. From this, we can deduce the correct relaxation of I in BV: $I_3(u) := \inf_{(\beta_0, \beta_1) \in \mathbb{R}^2} I_2(u, (\beta_0, \beta_1))$. However, this is far less natural than I_2 , and essential information encoded by β_1 at $x = 1$ is lost that way.

The example can be easily generalized to higher dimensions by considering $\Omega \subset \mathbb{R}^N$ a bounded domain of class C^1 with two disjoint parts of the boundary Γ_0 and Γ_1 of positive $N - 1$ dimensional Hausdorff measure. For instance, for $u \in W^{1,1}(\Omega; \mathbb{R}^M)$ and

$u(0) = 0$ on Γ_0 , a rough analogue of I (with a slightly modified boundary term to ensure $J < \infty$ on $W^{1,1}$) is given by

$$\text{minimize } J(u) := \int_{\Omega} (\text{dist}^2(x, \Gamma_1) + \varepsilon) |\nabla u(x)| \, dx + \int_{\Gamma_1} \sqrt{1 + (u(x) - \bar{u}(x))^2} \, d\mathcal{H}^{N-1}(x),$$

where $\Omega \subset \mathbb{R}^N$ is a bounded domain with a C^1 boundary, $\Gamma_1 \subset \partial\Omega$, and \bar{u} is a given function in $L^1(\Gamma_1, \mathcal{H}^{N-1} \llcorner \partial\Omega; \mathbb{R}^M)$. Above, “dist” denotes the distance function.

Concerning relaxation of this kind of functional, a natural replacement for the space X above is the Souček space $W_{\mu}^1(\bar{\Omega}) \cong \text{BV}(\Omega) \times \mathcal{M}(\partial\Omega)$, keeping track of the “true” boundary value as a measure on $\partial\Omega$ (the so-called outer trace in $W_{\mu}^1(\bar{\Omega})$). For more details on $W_{\mu}^1(\bar{\Omega})$ we refer to section 7.

The previous example shows that concentration effects at the boundary can be quite relevant for minimization problems with linear growth at infinity, even if all terms of the functional are convex, non-negative, and coercive. Once oscillations favored by nonconvex integrands come into play on top of that, we are lead to study the generalized gradient Young measures appearing as the natural limiting object capturing this behavior.

The main goal of our work is to complete the characterization of generalized Young measures given in [20] by deriving necessary and sufficient conditions which must be satisfied at the boundary of the domain Ω . In particular, our main result, summarized in Theorem 3.1, allows us to easily set up and prove relaxation results for a large class problems like our example, first in the space of generalized gradient Young measures and then in the Souček space, as a straightforward corollary.

The plan of the paper is as follows. We first introduce necessary notation and collect some auxiliary results in section 2. Then we state the main result, i.e., characterization of generalized gradient measures up to the boundary; cf. section 3. Proofs of necessity and sufficiency of our new condition at the boundary are the subject of sections 5 and 6, respectively. The first one, in section 7, is a relationship of generalized Young measures to the Souček space [30] which is a kind of a finer completion of $W^{1,1}(\Omega; \mathbb{R}^M)$ than $\text{BV}(\Omega; \mathbb{R}^M)$. As an application of our results, we then present a relaxation theorem for functionals on $W^{1,1}(\Omega; \mathbb{R}^M)$ including integral terms on the boundary in section 9. Our work then closes with the Appendix, which has three sections. Its first section contains a proof of an auxiliary result related to weak* lower semicontinuity along sequences of gradients which do not oscillate but concentrate at the boundary. The second section states a relationship between generalized Young measures and DiPerna–Majda measures often used by the second and the third author to characterize oscillations/concentrations in sequences of gradients. Finally, in the third section we explain the connection between quasiconvexity at the boundary and a variant of it which appears as an assumption in our relaxation theorem.

2. Notations and preliminaries. We start with some definitions and explanations of frequently used notation.

2.1. Basic notation. Throughout the text, unless otherwise specified, we will use the following notation.

- $\Omega \subset \mathbb{R}^N$, $N \geq 1$, stands for a bounded domain with a boundary of class C^1 .
- $|K|$ denotes the Lebesgue measure of $K \subset \mathbb{R}^N$.
- B denotes the open unit ball in \mathbb{R}^N centered at zero.
- Given a unit vector $\varrho \in \mathbb{R}^N$, we define the unit half-ball

$$(2.1) \quad D_{\varrho} = B \cap \{x \in \mathbb{R}^N; \varrho \cdot x < 0\}$$

- and $\Gamma_\varrho := \{x \in \partial D_\varrho; \varrho \cdot x = 0\} \subset \partial D_\varrho$.
- $B_r(x) := B(x, r) := \{y : |y - x| < r\}$ ($r > 0$ and the center point x is in a Euclidean vector space, with norm $|\cdot|$, that should be clear from the context).
- $(K)_r := \bigcup_{x \in K} B_r(x)$ (open r -neighborhood of a set $K \subset \mathbb{R}^N$).
- The unit spheres centered at zero in \mathbb{R}^N and $\mathbb{R}^{M \times N}$ are denoted by S^{N-1} and S^{MN-1} , respectively.
- Given $a \in \mathbb{R}^M$ and $b \in \mathbb{R}^N$, $a \otimes b$ is the matrix given by $(a \otimes b)_{ij} = a_i b_j$ for all $1 \leq i \leq M$ and all $1 \leq j \leq N$.
- Given real-valued functions $x \mapsto f(x)$ and $y \mapsto g(y)$, the function $(x, y) \mapsto (f \otimes g)(x, y)$ is defined by $(f \otimes g)(x, y) := f(x)g(y)$.

2.2. Spaces of functions and measures. By $\mathcal{M}(X)$ we denote the set of finite Radon measures on a Borel set X and by $\mathcal{M}_1^+(X)$ its subset of probability measures. We recall that the support of a measure μ , $\text{supp } \mu$, is the smallest closed set X such that $\mu(A) = 0$ if $X \cap A = \emptyset$. Given a measure μ , we write “ μ -almost all” or “ μ -a.e.” if we mean “up to a set with the μ -measure zero.” If $\mu = \mathcal{L}^N$, the N -dimensional Lebesgue measure, we usually omit writing it in the notation. As usual \mathcal{H}^{N-1} represents the $(N - 1)$ -dimensional Hausdorff measure in \mathbb{R}^N .

Given a domain $\Omega \subset \mathbb{R}^N$, the space $C(\Omega)$ consists of all real-valued continuous functions defined in Ω . Its subset $C_0(\Omega)$ is the completion with respect to the supremum norm of the space of functions with compact support in Ω ; that is, it consists of those functions $u \in C(\Omega)$ vanishing at the boundary ($u(x) \rightarrow 0$ as $\text{dist}(x; \partial\Omega) \rightarrow 0$) and at infinity ($u(x) \rightarrow 0$ as $|x| \rightarrow \infty$), in case the domain is unbounded. Recall that $C_0(\Omega)$ is a Banach space with respect to the norm $\|u\|_{C_0(\Omega)} = \sup_{x \in \Omega} |u(x)|$. $W^{1,p}(\Omega; \mathbb{R}^m)$, $1 \leq p \leq +\infty$, is the usual Sobolev space of measurable mappings which together with their first (distributional) derivatives are p th-integrable (if $p < +\infty$) or essentially bounded (if $p = +\infty$). The space of functions with bounded variation, $\text{BV}(\Omega; \mathbb{R}^M)$ (see [2]), consists of all $u \in L^1(\Omega; \mathbb{R}^M)$ such that its first order distributional derivatives $D_j u_i \in \mathcal{M}(\Omega)$. Given $u \in \text{BV}(\Omega; \mathbb{R}^M)$, we denote by Du the matrix-valued measure whose entries are $D_j u_i$, ∇u is the density of the absolutely continuous part of Du with respect to the Lebesgue measure, while Du^s denotes the singular part of Du with polar decomposition $dDu^s(x) = \frac{dDu^s}{d|Du^s|}(x) d|Du^s|(x)$. We recall that $(u_n)_{n \in \mathbb{N}} \subset \text{BV}(\Omega; \mathbb{R}^M)$ converges weakly* to $u \in \text{BV}(\Omega; \mathbb{R}^M)$ if $u_n \rightarrow u$ strongly in $L^1(\Omega; \mathbb{R}^M)$ and $Du_n \xrightarrow{*} Du$ in $\mathcal{M}(\Omega; \mathbb{R}^{M \times N})$; see [2, Definition 3.11].

Given $\mu \in \mathcal{M}(\Omega)$, we denote by $|\mu|$ its total variation (norm), i.e., $|\mu| := \sup \left| \int_\Omega f \, d\mu \right|$ where the supremum is taken over all $f \in C_0(\Omega)$ such that $\|f\|_{C_0(\Omega)} = 1$.

2.3. Quasiconvex functions and functions with linear growth. Let $\Omega \subset \mathbb{R}^N$ be a bounded Lipschitz domain. A function $v : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ is said to be quasiconvex [9] if for any $A \in \mathbb{R}^{M \times N}$ and any $\varphi \in W_0^{1,\infty}(\Omega; \mathbb{R}^M)$

$$(2.2) \quad v(A)|\Omega| \leq \int_\Omega v(A + \nabla\varphi(x)) \, dx.$$

If $v : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ is not quasiconvex, its quasiconvex envelope $Qv : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}$ is defined by

$$Qv = \sup \{ h \leq v; h : \mathbb{R}^{M \times N} \rightarrow \mathbb{R} \text{ quasiconvex} \}$$

or $Qv = -\infty$ if the set on the right-hand side is empty. If v is locally bounded and Borel measurable, then for any $A \in \mathbb{R}^{M \times N}$ (see [9])

$$(2.3) \quad Qv(A) = \inf_{\varphi \in W_0^{1,\infty}(\Omega; \mathbb{R}^M)} \frac{1}{|\Omega|} \int_\Omega v(A + \nabla\varphi(x)) \, dx.$$

Given a real-valued function v on some Euclidean space with norm $|\cdot|$, v is said to have *at most linear growth* if

$$|v(\cdot)| \leq C(1 + |\cdot|) \quad \text{for a constant } C \geq 0.$$

Clearly “at most linear growth” only refers to the behavior of v for large norms of its arguments.

Throughout this work, we use nonlinear transformations of measures, more precisely, derivatives in BV. A standard class of functions for which such nonlinear expressions can be defined is

$$(2.4) \quad \Upsilon := \{v \in C(\mathbb{R}^{M \times N}); v \text{ has at most linear growth \& } v^\infty \text{ exists}\},$$

where v^∞ denotes the *recession function* of v (if it exists). We recall that a function v is positively one-homogeneous if $v(\alpha \cdot) = \alpha v(\cdot)$ for all $\alpha \geq 0$ and that $v^\infty : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ is the (automatically continuous and positively one-homogeneous) function defined by

$$(2.5) \quad v^\infty(A) := \lim_{\alpha \rightarrow +\infty, t \rightarrow A} \frac{1}{\alpha} v(\alpha t).$$

If v^∞ exists, it coincides with the *generalized recession function* v^\sharp of v given by

$$(2.6) \quad v^\sharp(A) := \limsup_{\alpha \rightarrow +\infty} \frac{1}{\alpha} v(\alpha A).$$

Note, however, that even if v^\sharp is continuous and the limsup in (2.6) is a limit, this does not guarantee (2.5).

2.4. Tools and notation for sequences in BV. Here, we briefly recall some results of [5] which will be used below. In particular, we rely on the following notion:

DEFINITION 2.1. *Given a sequence $(u_n) \subset \text{BV}(\Omega; \mathbb{R}^M)$ and a closed set $K \subset \overline{\Omega}$, we say that (Du_n) does not charge K if $|Du_n|$ is tight in $\overline{\Omega} \setminus K$, i.e.,*

$$\sup_{n \in \mathbb{N}} |Du_n| \left((K)_r \cap \Omega \right) \xrightarrow{r \rightarrow 0^+} 0.$$

LEMMA 2.2 (Local decomposition in BV [5, Lemma 4.2]). *Let $\Omega \subset \mathbb{R}^N$ be open and bounded and let $K_j \subset \overline{\Omega}$, $j = 1, \dots, J$, be a finite family of compact sets such that $\overline{\Omega} \subset \bigcup_j K_j$. Then for every bounded sequence $(u_n) \subset \text{BV}(\Omega; \mathbb{R}^M)$ with $u_n \rightarrow 0$ in $L^1(\Omega; \mathbb{R}^M)$, there exists a subsequence (u_n) (not relabeled) which can be decomposed as*

$$u_n = u_{1,n} + \dots + u_{J,n},$$

where for each $j \in \{1, \dots, J\}$, $(u_{j,n})_n$ is a bounded sequence in $\text{BV}(\Omega; \mathbb{R}^M)$ converging to zero in L^1 such that the following two conditions hold for every j :

- (i) $\{u_{j,n} \neq 0\} \subset \{u_n \neq 0\}$, $\overline{\{u_{j,n} \neq 0\}} \subset (K_j)_{\frac{1}{n}} \setminus \bigcup_{i < j} (K_i)_{\frac{1}{2n}}$,
 $|Du_{j,n}|$ is absolutely continuous w.r.t. $|Du_n| + \mathcal{L}^N$,
 and $|Du_{j,n}| \leq |Du_n| + \frac{1}{n}$ as measures;
- (ii) $(Du_{j,n})$ does not charge $\bigcup_{i < j} K_i$.

Moreover, if $\partial\Omega$ is Lipschitz and each u_n has vanishing trace on $\partial\Omega$, this is inherited by $u_{j,n}$.

As derived in [5], the properties of the component sequences found in Lemma 2.2 guarantee that, at least asymptotically as $n \rightarrow \infty$, they behave as if they had pairwise disjoint support. To present this result we recall that if $v \in \Upsilon$ and $u \in \text{BV}(\Omega; \mathbb{R}^M)$, $v(Du)$ is defined as the real-valued measure given by

$$(2.7) \quad dv(Du)(x) := v(\nabla u(x))dx + v^\infty \left(\frac{dDu^s}{d|Du^s|}(x) \right) d|Du^s|(x),$$

where Du^s is the singular part of Du w.r.t. \mathcal{L}^N and ∇u is the density of the absolutely continuous part.

PROPOSITION 2.3 (cf. [5, Proposition 4.3]). *Suppose that $v \in \Upsilon$. Then for every $u \in \text{BV}(\Omega; \mathbb{R}^M)$ and every decomposition of a bounded sequence $u_n = u_{1,n} + \dots + u_{J,n} \in \text{BV}(\Omega; \mathbb{R}^M)$ with the properties listed in Lemma 2.2, we have that*

$$v(Du_n + Du) - v(Du) - \sum_{j=1}^J [v(Du_{j,n} + Du) - v(Du)] \xrightarrow[n \rightarrow \infty]{} 0$$

in total variation of measures.

Another, closely related observation is that for sequences purely concentrating at the boundary (but not in the interior), it is always possible to add or remove a nonzero weak* limit:

PROPOSITION 2.4 (cf. [5, Proposition 5.4] and its proof). *Suppose that $v \in \Upsilon$, and let $(c_n)_{n \in \mathbb{N}} \subset \text{BV}(\Omega; \mathbb{R}^M)$ be a bounded sequence such that $S_n := \{c_n \neq 0\} \subset (\partial\Omega)_{r_n}$ with a decreasing sequence $r_n \searrow 0$. Then for every $u \in \text{BV}(\Omega; \mathbb{R}^M)$,*

$$v(Du + Dc_n) - v(Du) - v(Dc_n) + v(0) \xrightarrow[n \rightarrow \infty]{} 0$$

in total variation of measures.

2.5. Functions with quasi-sublinear growth from below. This is a central condition for our analysis and it first appeared in [5].

DEFINITION 2.5 (Quasi-sublinear growth from below). *Assume that $x_0 \in \partial\Omega$. We say that a function $v : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ is quasi-sublinear from below (qslb) at x_0 (in short, $v \in \text{QSLB}(x_0)$) if*

for every $\varepsilon > 0$, there exist $\delta = \delta(\varepsilon) > 0$, $C = C(\varepsilon) \in \mathbb{R}$ s.t.

$$\int_{\Omega \cap B_\delta(x_0)} v(\nabla u(x)) dx \geq -\varepsilon \int_{\Omega \cap B_\delta(x_0)} |\nabla u(x)| dx - C$$

for every $u \in W^{1,1}(B_\delta(x_0) \cap \Omega; \mathbb{R}^M)$ with compact support in $B_\delta(x_0)$.

If $\partial\Omega$ is of class C^1 near $x_0 \in \partial\Omega$ and $v \in \Upsilon$ then we have the following equivalent variant of Definition 2.5: $v \in \text{QSLB}(x_0)$ if and only if for every $\varepsilon > 0$ there exists $C = C(\varepsilon) \geq 0$ such that

$$(2.8) \quad \int_{D_{\varrho(x_0)}} v(\nabla u(x)) dx \geq -\varepsilon \int_{D_{\varrho(x_0)}} |\nabla u(x)| dx - C$$

for all $u \in W_0^{1,1}(B; \mathbb{R}^M)$, where $\varrho(x_0)$ is the outer unit normal to $\partial\Omega$ at x_0 and $D_{\varrho(x_0)}$ is the half-ball as defined in (2.1).

In this case, we write (with a slight abuse of the notation) $v \in \text{QSLB}(\varrho(x_0))$ to indicate the dependence of the property of v on the normal vector to the boundary. Definition 2.5 corresponds to the condition used in Theorem 1.6 (ii) in [21] for $p = 1$ and to the notion of 1-quasi-subcritical growth from below at a boundary point as defined in [22].

Remark 2.6.

- (i) As the name suggests, quasi-sublinear growth from below is only affected by the behavior of v for large arguments. It holds that $v \in \text{QSLB}(x_0)$ if and only if $v^\infty \in \text{QSLB}(x_0)$, assuming that v^∞ (see (2.5)) exists. Moreover, for positively one-homogeneous functions such as v^∞ , (2.8) can be simplified significantly. More precisely, we have $v^\infty \in \text{QSLB}(\varrho(x_0))$ if and only if

$$(2.9) \quad \int_{D_{\varrho(x_0)}} v^\infty(\nabla u(x)) \, dx \geq 0$$

for all $u \in W_0^{1,1}(B; \mathbb{R}^M)$. For more details, we refer to [5, section 3].

- (ii) A “classical” counterpart of quasi-sublinear growth from below can be obtained in the same way one can widen the class of test functions in the definition of quasiconvexity to obtain a stronger condition equivalent to convexity, i.e., by removing the requirement that the test functions used under the integral are gradients. To do that in (2.8), we have to replace ∇u (with $u \in W_0^{1,1}(B; \mathbb{R}^M)$) by the larger class of test functions

$$U \in L^1(D_{\varrho(x_0)}; \mathbb{R}^{M \times N}) \quad \text{with} \quad \int_{D_{\varrho(x_0)}} U(x)\tau \, dx = 0 \quad \text{for all } \tau \in \mathbb{R}^N$$

$$\text{s.t. } \varrho(x_0) \cdot \tau = 0.$$

This formal link motivates the name for property (2.8), because (among other things) we may consider $U(x) = a \otimes \varrho(x_0)$ with an arbitrary constant vector $a \in \mathbb{R}^M$, and for such test functions instead of ∇u , (2.8) becomes equivalent to

$$\forall \varepsilon > 0 \exists C \geq 0 : v(a \otimes \varrho(x_0)) \geq -\varepsilon |a \otimes \varrho(x_0)| - C.$$

This in turn is equivalent to

$$0 \leq \liminf_{|a| \rightarrow +\infty, a \in \mathbb{R}^M} \frac{v(a \otimes \varrho(x_0))}{|a|},$$

i.e., sublinear growth of negative part of v , at least in all the “normal” directions $a \otimes \varrho(x_0)$. However, using all admissible U , the condition in fact also implies convexity at zero of the recession function v^∞ in the “tangential” directions $a \otimes \tau$, $\tau \cdot \varrho(x_0) = 0$.

- (iii) Unlike quasiconvexity, neither (2.8) nor its simpler version (2.9) for recession functions are Jensen-type inequalities. This is due to the lack of conditions on u on the flat part of $\partial D_{\varrho(x_0)}$, whereby the average of Du on $D_{\varrho(x_0)}$ is not fully controlled—it can be any matrix of the form $a \otimes \varrho(x_0)$, $a \in \mathbb{R}^M$. Interpreted from the point of view of convex analysis, (2.9) states that zero is an element of a kind of generalized subdifferential (in the “quasi-at-the-boundary” sense) of v^∞ at zero. This is more restrictive than quasiconvexity at the boundary at zero, because for the latter a nonempty generalized subdifferential at zero is enough (see (C.1) at $A = 0$ and (C.2) in Appendix C). In fact, there are even

linear functions that do not satisfy (2.9), for instance $v^\infty(A) := (b \otimes \varrho(x_0)) : A$ for any $b \in \mathbb{R}^M \setminus \{0\}$, due to Lemma 2.10 below (with $a := -b$). Notice that this example features linear growth of the negative part of v^∞ in the “normal” direction $-b \otimes \varrho(x_0)$.

Functions with quasi-sublinear growth from below play a key role in characterization of weak* lower semicontinuity in BV. We recall the following definition.

DEFINITION 2.7 (w*-lsc). *We say that a functional $F : \text{BV}(\Omega; \mathbb{R}^M) \rightarrow \mathbb{R}$ is sequentially weak* lower semicontinuous (w*-lsc) in $\text{BV}(\Omega; \mathbb{R}^M)$ if*

$$\liminf_{n \rightarrow \infty} F(u_n) \geq F(u)$$

for every sequence $(u_n) \subset \text{BV}(\Omega; \mathbb{R}^M)$ and such that $u_n \overset{*}{\rightharpoonup} u$ in BV.

Let $f \in C(\bar{\Omega}; \Upsilon)$ and $F : \text{BV}(\Omega; \mathbb{R}^M) \rightarrow \mathbb{R}$ be defined as

$$(2.10) \quad F(u) := \int_{\Omega} df(x, Du)(x),$$

where

$$df(x, Du) := f(x, \nabla u(x))dx + f^\infty \left(x, \frac{dDu}{d|Du|}(x) \right) d|D^s u|(x).$$

The proof of next proposition, Proposition 2.8, is implicitly contained in that of Theorem 2.9 in [5]. It shows that being of quasi-sublinear growth from below is sufficient to ensure weak* lower semicontinuity along sequences concentrating at the boundary. For the convenience of the reader, a proof of this proposition can be found in Appendix A.

PROPOSITION 2.8. *Assume that $f = g \otimes v$ with $(g, v) \in C(\bar{\Omega}) \times \Upsilon$, $f(x, \cdot) \in \text{QSLB}(x)$ for all $x \in \partial\Omega$ and F is given by (2.10). Then F is weak* lower semicontinuous along all sequences $(c_n)_{n \in \mathbb{N}} \subset \text{BV}(\Omega; \mathbb{R}^N)$ that are bounded and satisfy $S_n := \{c_n \neq 0\} \cup \text{supp} |Dc_n| \subset (\partial\Omega)_{r_n}$ for a decreasing sequence $r_n \searrow 0$.*

The dependence on the normal at a given boundary point is illustrated by the following lemma.

LEMMA 2.9. *Let $v : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ be continuous and with linear growth, and let $\varrho_1, \varrho_2 \in S^{N-1}$. Then we have the following:*

- (i) *If $R_{21} \in \mathbb{R}^{N \times N}$ is an orthogonal matrix such that $\varrho_2 = R_{21}\varrho_1$, then $v \in \text{QSLB}(\varrho_1)$ if and only if $A \mapsto v(AR_{21}) \in \text{QSLB}(\varrho_2)$.*
- (ii) *Suppose in addition that v has a recession function v^∞ in the sense of (2.5). Then for each $v \in \text{QSLB}(\varrho_1)$ and every $\alpha > 0$, there exists $\beta > 0$ such that $A \mapsto v(A) + \alpha|A| \in \text{QSLB}(\varrho_2)$ for every $\varrho_2 \in S^{N-1}$ satisfying $|\varrho_1 - \varrho_2| < \beta$.*

Proof.

(i): Let $\varphi_1 \in W_0^{1,1}(B; \mathbb{R}^M)$. We have that

$$\int_{D_{e_1}} v(\nabla \varphi_1) dx \geq -\varepsilon \int_{D_{e_1}} |\nabla \varphi_1| dx - C_\varepsilon$$

if and only if for $\varphi_2 \in W_0^{1,1}(B_1; \mathbb{R}^M)$ and $\varphi_2(y) := \varphi_1(R_{21}^{-1}y)$,

$$\int_{D_{e_2}} v((\nabla \varphi_2)R_{21}) dy \geq -\varepsilon \int_{D_{e_2}} |\nabla \varphi_2| dy - C_\varepsilon,$$

by the change of variables given by $y = R_{21}x$. Here, note that $D_{\varrho_2} = R_{21}D_{\varrho_1}$, $|\det R_{21}| = 1$, and $|(\nabla\varphi_2)R_{21}| = |\nabla\varphi_2|$.

- (ii): For each $\varrho_2 \in S^{N-1}$, choose an orthogonal $R_{21} \in \mathbb{R}^{N \times N}$ such that $\varrho_2 = R_{21}\varrho_1$ and $|I - R_{21}| = |\varrho_1 - \varrho_2| < \beta$, where $I \in \mathbb{R}^{N \times N}$ is the identity matrix. (For convenience and to avoid additional constants, we are assuming here that the norm $|\cdot|$ used in $\mathbb{R}^{N \times N}$ is the operator norm.) By Remark 2.6 and the fact that $(v + \alpha|\cdot|)^\infty = v^\infty + \alpha|\cdot|$, it suffices to show the assertion for $v = v^\infty$. Since v^∞ is one-homogeneous and uniformly continuous on the unit sphere, there exists a nondecreasing continuous function $\omega : [0, \infty) \rightarrow [0, \infty)$ with $\omega(0) = 0$ such that for all orthogonal $R \in \mathbb{R}^{N \times N}$,

$$|v^\infty(A) - v^\infty(AR)| \leq |A|\omega(|I - R|).$$

As a consequence, we can choose $\beta > 0$ small enough so that whenever $|\varrho_1 - \varrho_2| < \beta$ (and therefore $|I - R_{21}| < \beta$),

$$|v^\infty(A) - v^\infty(AR_{21})| \leq |A|\omega(|I - R_{21}|) \leq \alpha|A| \quad \text{for every } A \in \mathbb{R}^{M \times N}.$$

Using the characterization of QSLB for recession functions given in (2.9) and the fact that $v^\infty(\cdot R_{21}) \in \text{QSLB}(\varrho_2)$ according to (i), we infer that for every $\varphi \in W_0^{1,1}(B; \mathbb{R}^M)$,

$$\int_{D_{\varrho_1}} (v^\infty(\nabla\varphi) + \alpha|\nabla\varphi|) dx \geq \int_{D_{\varrho_1}} v^\infty((\nabla\varphi)R_{21}) dx \geq 0.$$

Therefore, $v^\infty + \alpha|\cdot| \in \text{QSLB}(\varrho_2)$. □

Similar to the well-known relationship of quasiconvexity and rank-1-convexity, the following can be said about functions in QSLB:

LEMMA 2.10. *Given $\varrho \in S^{N-1}$, let $v \in \text{QSLB}(\varrho)$ and be positively one-homogeneous. Then for all matrices of the form $A := a \otimes \varrho$ with arbitrary $a \in \mathbb{R}^M$, we have that $v(A) \geq 0$.*

Proof. The proof is indirect. Assume by contradiction that $v(A) < 0$ for an admissible matrix A . Let $\varepsilon, r \in (0, \frac{1}{2})$ and define

$$S(r) := \{y \in \partial D_\varrho, |y| < r\} \subset \partial D_\varrho$$

(see (2.1)) and let

$$\begin{aligned} Z^-(r, \varepsilon) &:= \{y - t\varrho \mid y \in \partial D_\varrho, |y| < r, 0 < t < \varepsilon\} \subset D_\varrho, \\ Z^+(r, \varepsilon) &:= \{y + t\varrho \mid y \in \partial D_\varrho, |y| < r, 0 < t < \varepsilon\} \subset -D_\varrho \end{aligned}$$

denote the two cylinders of height ε on either side of their circular base $S(r)$ located in the flat part of ∂D_ϱ . Finally, let $Z(r, \varepsilon)$ denote the interior of $Z^-(r, \varepsilon) \cup Z^+(r, \varepsilon)$. For fixed $r > 0$, we choose $\varphi_\varepsilon \in W_0^{1,\infty}(B)$ in such a way that

$$\nabla\varphi_\varepsilon = A \text{ in } Z^-(r, \varepsilon), \quad \nabla\varphi_\varepsilon = -A \text{ in } Z^+(r, \varepsilon), \quad \nabla\varphi_\varepsilon = 0 \text{ in } B \setminus \overline{Z(r + \varepsilon, \varepsilon)},$$

and

$$|\nabla\varphi_\varepsilon| \leq |A| \text{ in } Z(r + \varepsilon, \varepsilon) \setminus \overline{Z(r, \varepsilon)},$$

which is, basically, (for $r \gg \varepsilon$) a long, narrow “tent” on $Z(r, \varepsilon)$ closed with “face walls” of similar slope above $Z(r + \varepsilon, \varepsilon) \setminus \overline{Z(r, \varepsilon)}$. Due to $v \in \text{QSLB}(\varrho)$, we have

$$\begin{aligned} 0 &\leq \frac{1}{\varepsilon} \int_{D_\varepsilon} v(\nabla \varphi_\varepsilon) \, dx \\ &= \frac{1}{\varepsilon} |Z^-(r, \varepsilon)| v(A) + \frac{1}{\varepsilon} \int_{Z^-(r+\varepsilon, \varepsilon) \setminus \overline{Z^-(r, \varepsilon)}} v(\nabla \varphi_\varepsilon) \, dx \\ &\leq \mathcal{H}^{N-1}(S(r))v(A) + \mathcal{H}^{N-1}(S(r + \varepsilon) \setminus S(r)) |A| \|v\|_{L^\infty(S^{M \times N-1})}. \end{aligned}$$

Since $v(A) < 0$ and $\mathcal{H}^{N-1}(S(r + \varepsilon) \setminus S(r)) \leq C\varepsilon$, we obtain a contradiction in the limit as $\varepsilon \rightarrow 0$. □

Remark 2.11. For general functions $v \in \text{QSLB}(\varrho)$ possessing a recession function v^∞ , we automatically have $v^\infty \in \text{QSLB}(\varrho)$, and by continuity, one-homogeneity and (2.5),

$$v^\infty(A) = \lim_{t \rightarrow +\infty} \frac{1}{t} v^\infty(\xi + tA) = \lim_{t \rightarrow +\infty} \frac{1}{t} v(\xi + tA)$$

for all $\xi, A \in \mathbb{R}^{M \times N}$. For $v \in \text{QSLB}(\varrho)$, the assertion of Lemma 2.10 therefore generalizes to

$$\limsup_{t \rightarrow +\infty} \frac{1}{t} v(\xi + t(a \otimes \varrho)) \geq 0 \quad \text{for all } \xi \in \mathbb{R}^{M \times N}, a \in \mathbb{R}^M,$$

i.e., v has sublinear growth from below along rank-1 lines corresponding to the normal ϱ .

2.6. Young measures and their generalizations.

2.6.1. Young measures: The sets $\mathcal{Y}(\Omega; \mathbb{R}^{M \times N})$ and $\mathcal{Y}^1(\Omega; \mathbb{R}^{M \times N})$. *Young measures* on a bounded domain $\Omega \subset \mathbb{R}^N$ are weakly*-measurable mappings $x \mapsto \nu_x, x \in \Omega$, with $\nu_x : \Omega \rightarrow \mathcal{M}_1^+(\mathbb{R}^{M \times N})$. We recall that the adjective “weakly*-measurable” means that, for any $v \in C_0(\mathbb{R}^{M \times N})$, the mapping $x \mapsto \int_{\mathbb{R}^{M \times N}} v(A) d\nu_x(A), x \in \Omega$, is measurable in the usual sense. Let us also recall here that, by the Riesz theorem, $\mathcal{M}(\mathbb{R}^{M \times N})$, normed by the total variation, is a Banach space which is isometrically isomorphic with $C_0(\mathbb{R}^{M \times N})^*$ (the dual space of $C_0(\mathbb{R}^{M \times N})$).

We denote the set of all Young measures by $\mathcal{Y}(\Omega; \mathbb{R}^{M \times N})$. It is known that $\mathcal{Y}(\Omega; \mathbb{R}^{M \times N})$ is a convex subset of $L_{w^*}^\infty(\Omega; \mathcal{M}(\mathbb{R}^{M \times N})) \cong L^1(\Omega; C_0(\mathbb{R}^{M \times N}))^*$, where the subscript “w*” indicates the property “weakly*-measurable.”

Let

$$(2.11) \quad C_1(\mathbb{R}^{M \times N}) = \left\{ v \in C(\mathbb{R}^{M \times N}); \lim_{|A| \rightarrow +\infty} v(A)/|A| = 0 \right\}.$$

As it was shown in [3, 32, 33], for every bounded sequence $(Y_k)_{k \in \mathbb{N}} \subset L^1(\Omega; \mathbb{R}^{M \times N})$, there exists a subsequence (not relabeled) and a Young measure $\nu = \{\nu_x\}_{x \in \Omega} \in \mathcal{Y}(\Omega; \mathbb{R}^{M \times N})$ such that for all $v \in C_1(\mathbb{R}^{M \times N})$ (see (2.11)) and all $g \in L^\infty(\Omega)$

$$(2.12) \quad \lim_{k \rightarrow \infty} \int_\Omega g(x) v(Y_k(x)) \, dx = \int_\Omega g(x) \left[\int_{\mathbb{R}^{M \times N}} v(A) d\nu_x(A) \right] \, dx.$$

Let us denote by $\mathcal{Y}^1(\Omega; \mathbb{R}^{M \times N})$ the set of all Young measures which are created in this way, i.e., by considering all bounded sequences in $L^1(\Omega; \mathbb{R}^{M \times N})$.

If v has linear growth at infinity, the limit passage (2.12) generally does not hold due to concentration effects created by (Y_k) if this sequence is not uniformly integrable. For this reason, various generalizations have been invented to allow for a description of limits in (2.12). We refer to [29] for a survey of these approaches. The authors in [1] showed that if $(Y_k)_{k \in \mathbb{N}} \subset L^1(\Omega; \mathbb{R}^{M \times N})$ is a bounded sequence then there is a (non-re-labeled) subsequence, a Young measure $\nu \in \mathcal{Y}^1(\Omega; \mathbb{R}^{M \times N})$, a positive measure $\lambda \in \mathcal{M}(\bar{\Omega})$, and a mapping $\nu^\infty \in L^\infty_{w*}(\bar{\Omega}, \lambda; \mathcal{M}(S^{M \times N-1}))$ with values in probability measures and defined for λ -almost all $x \in \bar{\Omega}$ such that for all $g \in C(\bar{\Omega})$ and all $v \in \Upsilon$ (cf. (2.4))

$$(2.13) \quad \lim_{k \rightarrow \infty} \int_{\Omega} g(x)v(Y_k(x)) \, dx = \int_{\Omega} g(x) \left[\int_{\mathbb{R}^{M \times N}} v(A) d\nu_x(A) \right] dx + \int_{\bar{\Omega}} g(x) \left[\int_{S^{M \times N-1}} v^\infty(A) d\nu_x^\infty(A) \right] d\lambda(x).$$

Roughly speaking, λ measures how much $(|Y_k|)$ concentrates and ν^∞ accounts for the spatial distribution of concentrations. A closely related description can also be found in [11] and [13].

Continuous functions possessing recession function can be continuously extended by radial limits to the compactification of $\mathbb{R}^{M \times N}$ by the sphere. Indeed, defining $d : \mathbb{R}^{M \times N} \rightarrow B$, $d(A) := A/(1+|A|)$ we can identify the compactification by the sphere with \bar{B} . Conversely, to any continuous function on \bar{B} we can assign a function on $\mathbb{R}^{M \times N}$ which has the recession function. Notice also that positively one-homogeneous functions are fully determined by their values on $S^{M \times N-1}$. If $v \in \Upsilon$ then we can assume that

$$\frac{v(A)}{1+|A|} = \begin{cases} c + v_{0,0}(A) + v_{0,1} \left(\frac{A}{|A|} \right) \frac{|A|}{1+|A|} & \text{if } A \neq 0, \\ c + v_{0,0}(0) & \text{if } A = 0, \end{cases}$$

where $v_{0,0} \in C_0(\mathbb{R}^{M \times N})$, $v_{0,1} \in C(S^{M \times N-1})$, and $c \in \mathbb{R}$. Consequently, for $g \in C(\bar{\Omega})$ the map $(x, s) \mapsto g(x)v(s)/(1+|s|) \in C(\bar{\Omega}) \otimes C(\bar{B})$. By the Stone-Weierstrass theorem equality (2.13) also holds for test functions of the form $(x, A) \mapsto f(x, A) = f_0(x, A/(1+|A|))(1+|A|)$, where $f_0 \in C(\bar{\Omega} \times \bar{B})$. Notice that such a function f always has a recession function given by $f^\infty(x, A) = |A| f_0(x, \frac{A}{|A|})$, satisfying an even stronger version of (2.5) additionally involving a kind of uniform convergence in x , namely,

$$(2.14) \quad f^\infty(x, A) = \lim_{\alpha \rightarrow +\infty, t \rightarrow A, y \rightarrow x} \frac{1}{\alpha} f(y, \alpha t).$$

In particular, f^∞ is always continuous in x . In general, (2.13) cannot be expected to hold with test functions f having a (generalized) recession function with discontinuities in x .

2.6.2. The set of generalized Young measures $\mathcal{Y}_{\text{gen}}^1(\bar{\Omega}; \mathbb{R}^{M \times N})$. We denote by $\Lambda := (\nu, \lambda, \nu^\infty)$ the triple appearing in (2.13), for some bounded sequence $(Y_k) \subset L^1(\Omega; \mathbb{R}^{M \times N})$, and the set of all such triples is represented by $\mathcal{Y}_{\text{gen}}^1(\bar{\Omega}; \mathbb{R}^{M \times N})$, or briefly $\mathcal{Y}_{\text{gen}}^1$. For the right-hand side of (2.13), given $g \in C(\bar{\Omega})$ and all $v \in \Upsilon$, we occasionally use the shorthand notation

$$(2.15) \quad \langle \langle \Lambda, g \otimes v \rangle \rangle := \int_{\Omega} g(x) \langle \nu_x, v \rangle dx + \int_{\bar{\Omega}} g(x) \langle \nu_x^\infty, v^\infty \rangle d\lambda(x),$$

where

$$\langle \nu_x, v \rangle := \int_{\mathbb{R}^{M \times N}} v(A) d\nu_x(A) \quad \text{and} \quad \langle \nu_x^\infty, v^\infty \rangle := \int_{S^{M \times N-1}} v^\infty(A) d\nu_x^\infty(A).$$

Also note that actually ν_x is only defined for a.e. $x \in \Omega$ (or, equivalently, a.e. $x \in \bar{\Omega}$, since $\partial\Omega$ has vanishing Lebesgue measure), and ν_x^∞ is only defined for λ -a.e. $x \in \bar{\Omega}$. Accordingly, given $\Lambda_1, \Lambda_2 \in \mathcal{Y}_{\text{gen}}^1$, $\Lambda_1 = (\nu_1, \lambda_1, \nu_1^\infty)$, and $\Lambda_2 = (\nu_2, \lambda_2, \nu_2^\infty)$, with a slight abuse of notation we write $\Lambda_1 = \Lambda_2$ if $\nu_{1,x} = \nu_{2,x}$ for a.e. $x \in \Omega$, $\lambda_1 = \lambda_2$ and $\nu_{1,x}^\infty = \nu_{2,x}^\infty$ for λ_1 or λ_2 -a.e. $x \in \bar{\Omega}$.

DEFINITION 2.12 (Generation). *We say that $\Lambda = (\nu, \lambda, \nu^\infty)$ is generated by a bounded sequence $(Y_k) \subset L^1(\Omega; \mathbb{R}^{M \times N})$ whenever (2.13) holds.*

We observe that this notion also makes sense for slightly more general sequences, namely, if (Y_k) is a sequence of $\mathbb{R}^{M \times N}$ -valued Radon measures on Ω with bounded total variation. In that case, we say that $\Lambda = (\nu, \lambda, \nu^\infty)$ is generated by (Y_k) if

$$(2.16) \quad \lim_{k \rightarrow \infty} \int_{\bar{\Omega}} g(x) dv(Y_k)(x) = \langle \langle \Lambda, g \otimes v \rangle \rangle$$

for every $g \in C(\bar{\Omega})$ and every $v \in \Upsilon$. Analogous to the special case of BV derivatives mentioned before, here

$$dv(Y_k)(x) := v \left(\frac{dY_k}{d\mathcal{L}^N}(x) \right) dx + v^\infty \left(\frac{dY_k}{d|Y_k|}(x) \right) d|Y_k^s|(x),$$

where $\frac{dY_k}{d\mathcal{L}^N}$ denotes the density of the absolutely continuous part of Y_k w.r.t. the Lebesgue measure, Y_k^s is its singular part, and $\frac{dY_k}{d|Y_k|}$ stands for the density of Y_k with respect to its total variation $|Y_k|$.

Example 2.13. Let $\Omega \subset \mathbb{R}^N$ a bounded domain with smooth boundary such that $0 \in \bar{\Omega}$, and take a function $Y \in L^1(B; \mathbb{R}^{M \times N})$ on the unit ball B centered at the origin, extended by zero to all of \mathbb{R}^N (without renaming it). We define $Y_k(x) := k^N Y(kx)$ for all $k \in \mathbb{N}$. Clearly $(Y_k)_{k \in \mathbb{N}}$ is uniformly bounded in $L^1(\Omega; \mathbb{R}^{M \times N})$ and $Y_k \rightarrow 0$ in measure. Consequently, (Y_k) generates $\Lambda \in \mathcal{Y}_{\text{gen}}^1$ such that $\nu_x = \delta_0$, $\lambda = \|Y\|_{L^1(D_\varrho; \mathbb{R}^{M \times N})} \delta_0$ (in case $0 \in \partial\Omega$), or $\lambda = \|Y\|_{L^1(B; \mathbb{R}^{M \times N})} \delta_0$ (in case $0 \in \Omega$), and

$$\begin{aligned} & \int_{S^{M \times N-1}} v^\infty(A) d\nu_0^\infty(A) \\ &= \begin{cases} \|Y\|_{L^1(D_\varrho; \mathbb{R}^{M \times N})}^{-1} \int_{D_\varrho} v^\infty(Y(z)) dz & \text{if } 0 \in \partial\Omega \text{ with outer normal } \varrho, \\ \|Y\|_{L^1(B; \mathbb{R}^{M \times N})}^{-1} \int_B v^\infty(Y(z)) dz & \text{if } 0 \in \Omega. \end{cases} \end{aligned}$$

Notice that in this particular example, $x = 0$ is the only relevant point for ν_x^∞ in $\bar{\Omega}$ because $\lambda(\bar{\Omega} \setminus \{0\}) = 0$. This kind of example is also possible for gradients, namely if $Y = \nabla u$ for some $u \in W^{1,1}(\mathbb{R}^N; \mathbb{R}^M)$. In this case, we let $Y_k = \nabla u_k$ with $u_k(x) := k^{N-1} u(kx)$, and consequently Λ is a generalized gradient Young measure as defined below.

2.6.3. Piecewise generation of generalized Young measures. The natural embedding of a function space containing a generating sequence, say, $L^1(\Omega; \mathbb{R}^{M \times N})$,

into $\mathcal{Y}_{\text{gen}}^1(\bar{\Omega}; \mathbb{R}^{M \times N})$, is not linear, because in $\mathcal{Y}_{\text{gen}}^1$ addition and multiplication with scalars is carried out *after* the application of a nonlinear test function v . Nevertheless, in some special cases, a kind of additivity can be observed. Essentially, we need two generating sequences lacking any kind of interaction. In terms of the generated generalized Young measures, this is made precise by requiring them to be orthogonal in the following measure theoretic sense:

DEFINITION 2.14. *Given $\Psi = (\eta_x, \mu, \eta_x^\infty), \Theta = (\vartheta_x, \sigma, \vartheta_x^\infty) \in \mathcal{Y}_{\text{gen}}^1$, we write that $\Psi \perp \Theta$ if there are two disjoint Borel sets $S, T \subset \bar{\Omega}$ such that $\bar{\Omega} = S \cup T$, $\mu(T) = 0 = \mathcal{L}^N(\{x \in T \mid \eta_x \neq \delta_0\})$, and $\sigma(S) = 0 = \mathcal{L}^N(\{x \in S \mid \vartheta_x \neq \delta_0\})$.*

We now claim that for such a pair the sum of the respective generating sequences will always generate the sum $\chi_S\Psi + \chi_T\Theta$. The key ingredient for the proof is the following uniform continuity property:

PROPOSITION 2.15 (Special case of [5, Proposition 4.5]). *Let $v \in \Upsilon$ and $g \in C(\bar{\Omega})$. Then for every pair of sequences $(\mu_k), (\lambda_k) \subset \mathcal{M}(\Omega; \mathbb{R}^{M \times N})$ such that $|\mu_k|(\Omega)$ and $|\lambda_k|(\Omega)$ are bounded,*

$$|\mu_k - \lambda_k| \rightarrow 0 \text{ implies that } \int_{\Omega} g(x)dv(\mu_k)(x) - \int_{\Omega} g(x)dv(\lambda_k)(x) \rightarrow 0.$$

LEMMA 2.16 (Additive behavior along generating sequences for orthogonal generalized Young measures). *Let $\Psi = (\eta_x, \mu, \eta_x^\infty), \Theta = (\vartheta_x, \sigma, \vartheta_x^\infty) \in \mathcal{Y}_{\text{gen}}^1$ be such that $\Psi \perp \Theta$, and let $\bar{\Omega} = S \cup T$ denote the associated decomposition of Ω of Definition 2.14. Moreover, let $(w_k), (z_k) \subset L^1$ be two bounded sequences generating Ψ and Θ , respectively. Then $(w_k + z_k)$ generates*

$$\chi_S\Psi + \chi_T\Theta := (\chi_S(x)\eta_x + \chi_T(x)\vartheta_x, \mu(S \cap \cdot) + \sigma(T \cap \cdot), \chi_S(x)\eta_x^\infty + \chi_T(x)\vartheta_x^\infty).$$

Remark 2.17. Lemma 2.16 also holds for more general generating sequences in \mathcal{M} instead of L^1 . The proof remains the same, apart from the fact that expressions like, say, $v(w_k)dx$ have to be replaced by $dv(w_k)(x)$.

Proof of Lemma 2.16. Let $\Lambda = (\nu_x, \lambda, \nu_x^\infty)$ be the generalized Young measure generated by $(w_k + z_k)$ (or a suitable subsequence, not relabeled). Since $(\Psi, S, (w_k))$ and $(\Theta, T, (z_k))$ are interchangeable, it suffices to show that $\Lambda = \Psi$ on S , i.e., $\nu_x = \eta_x$ for \mathcal{L}^N -a.e. $x \in S$, $\lambda|_S = \mu|_S$, and $\nu_x^\infty = \eta_x^\infty$ for λ -a.e. $x \in S$. That way, we obtain that for each subsequence of $(w_k + z_k)$ generating some generalized Young measure, that measure coincides with $\chi_S\Psi + \chi_T\Theta$, the same for all such subsequences, and consequently the whole sequence generates $\chi_S\Psi + \chi_T\Theta$.

By the inner regularity of the Borel measures $\tilde{\mu}, \tilde{\sigma}$ defined by

$$d\tilde{\mu}(x) := d\mu(x) + \langle \eta_x, \cdot \rangle dx, \quad d\tilde{\sigma}(x) := d\sigma(x) + \langle \vartheta_x, \cdot \rangle dx,$$

there exist increasing sequences of compact sets $S^{(m)} \subset S$ and $T^{(m)} \subset T$ such that for all $m \in \mathbb{N}$,

$$\left| \tilde{\mu}(S^{(m)}) - \tilde{\mu}(S) \right| \leq \frac{1}{m} \quad \text{and} \quad \left| \tilde{\sigma}(T^{(m)}) - \tilde{\sigma}(T) \right| \leq \frac{1}{m}.$$

Since $\tilde{\mu}(S \setminus \bigcup_{m \in \mathbb{N}} S^{(m)}) = 0$, it now suffices to show that $\Lambda = \Psi$ on $S^{(m)}$ for all m .

Suppose by contradiction that $\Lambda \neq \Psi$ on $S^{(m)}$ for an $m \in \mathbb{N}$. Consequently, there exist $\varepsilon > 0$, $g \in C(S^{(m)})$ and $v \in \Upsilon$ such that

$$(2.17) \quad \varepsilon < \left| \int_{S^{(m)}} g(x) (\langle \eta_x, v \rangle dx + \langle \eta_x^\infty, v^\infty \rangle d\mu(x)) - \int_{S^{(m)}} g(x) (\langle \nu_x, v \rangle dx + \langle \nu_x^\infty, v^\infty \rangle d\lambda(x)) \right|.$$

By definition of Υ , there is a constant $C > 0$ such that $|v(\cdot)| \leq C(1+|\cdot|)$ and $|v^\infty(\cdot)| \leq C|\cdot|$. Due to the outer regularity of Radon measures, there exists a neighborhood V of $S^{(m)}$ in $\bar{\Omega}$ such that

$$(2.18) \quad \|g\|_{L^\infty(S^{(m)})} \int_{V \setminus S^{(m)}} C((1 + \langle \bar{\nu}_x, |\cdot| \rangle)) dx + \langle \bar{\nu}_x^\infty, |\cdot| \rangle d\bar{\lambda}(x) \leq \frac{\varepsilon}{3},$$

where $\bar{\Lambda} := (\bar{\nu}_x, \bar{\lambda}, \bar{\nu}_x^\infty)$ denotes the generalized Young measure generated by $(|w_k| + |z_k|)$ (up to a subsequence). Observe that besides holding for $\bar{\Lambda}$, (2.18) also holds for any generalized Young measure generated by a sequence (y_k) such that $|y_k| \leq |w_k| + |z_k|$ a.e. in Ω . Without changing notation, we extend g to a function in $C(\bar{\Omega})$ in such a way that $g = 0$ on $\bar{\Omega} \setminus V$ and $\|g\|_{L^\infty(\bar{\Omega})} \leq \|g\|_{L^\infty(S^{(m)})}$.

On the other hand, for any function $\varphi \in C(\bar{\Omega})$, we have

$$\int_{\Omega} \varphi(x) |z_k| dx \rightarrow \langle \langle \Theta, \varphi \otimes |\cdot| \rangle \rangle = \int_{\bar{\Omega}} \varphi(x) d\tilde{\sigma}(x).$$

In particular, for arbitrary but fixed $m, h \in \mathbb{N}$ and any set $U = U(m, h) \subset \bar{\Omega}$ which is relatively open with respect to $\bar{\Omega}$ and satisfies $S^{(m)} \subset U \subset \bar{U} \subset \bar{\Omega} \setminus T^{(h)}$, we may choose $0 \leq \varphi \leq 1$ such that $\varphi = 1$ on U and $\varphi = 0$ on $T^{(h)}$, which yields that

$$(2.19) \quad \limsup_k \int_U |z_k| dx \leq \tilde{\sigma}(\bar{\Omega} \setminus T^{(h)}) = \tilde{\sigma}(T \setminus T^{(h)}) \leq \frac{1}{h}.$$

We claim that, as a consequence, we can choose $h \in \mathbb{N}$ large enough, together with an associated neighborhood $U = U(m, h)$ of $S^{(m)}$ in $\bar{\Omega}$ as introduced above, without loss of generality $U \subset V$, such that

$$(2.20) \quad \limsup_k \left| \int_U g(x)v(w_k + \chi_U z_k) dx - \int_U g(x)v(w_k) dx \right| \leq \frac{\varepsilon}{3}.$$

Indeed, if (2.20) failed for all large h , there would be a sequence $k(h) \rightarrow \infty$ as $h \rightarrow \infty$ such that

$$\liminf_h \left| \int_U g(x)v(w_{k(h)} + \chi_{U(m,h)} z_{k(h)}) dx - \int_U g(x)v(w_{k(h)}) dx \right| \geq \frac{\varepsilon}{3} > 0,$$

which is impossible due to Proposition 2.15 and (2.19).

Recall that (w_k) generates Ψ , and let Λ_U denote the generalized Young measure generated by $(w_k + \chi_U z_k)$ (up to a subsequence). Passing to the limit inside the modulus in (2.20), we get that

$$(2.21) \quad |\langle \langle \Lambda_U, g \otimes v \rangle \rangle - \langle \langle \Psi, g \otimes v \rangle \rangle| \leq \frac{\varepsilon}{3}.$$

As remarked above just below (2.18), besides holding for $\bar{\Lambda}$, (2.18) also holds for the generalized Young measures $\Lambda_U = (\nu_{U,x}, \lambda_U, \nu_{U,x}^\infty)$ and Ψ . Since $g = 0$ outside of V , we can apply these two estimates in (2.21), and infer that

$$(2.22) \quad \left| \int_{S^{(m)}} g(x) (\langle \nu_{U,x}, v \rangle dx + \langle \nu_{U,x}^\infty, v \rangle d\lambda_U(x)) - \int_{S^{(m)}} g(x) (\langle \eta_x, v \rangle dx + \langle \eta_x^\infty, v \rangle d\mu(x)) \right| \leq \varepsilon.$$

Since $S^{(m)}$ is compactly contained in U , the generating sequences of Λ and Λ_U coincide on a whole neighborhood of $S^{(m)}$, whence $\Lambda = \Lambda_U$ on $S^{(m)}$. Therefore, (2.22) contradicts (2.17). \square

2.6.4. Generalized gradient Young measures: $\text{GYM}(\bar{\Omega}; \mathbb{R}^M)$. The main topic of this article is an explicit characterization of the subclass of generalized Young measures generated by gradients.

DEFINITION 2.18 (Generalized gradient Young measures). *We say that $\Lambda := (\nu, \lambda, \nu^\infty) \in \mathcal{Y}_{\text{gen}}^1$ is a generalized BV-gradient Young measure if it is generated by a sequence of derivatives in BV, i.e., if (2.16) holds with $Y_k = Du_k$ for a bounded sequence $(u_k) \subset \text{BV}(\Omega; \mathbb{R}^M)$. The class of all generalized BV-gradient Young measures is denoted by $\text{GYM}(\bar{\Omega}; \mathbb{R}^M)$, or briefly GYM.*

Remark 2.19. As already pointed out in [20, Proposition 4], every $\Lambda \in \text{GYM}$ can be generated by (∇u_k) , for a suitable bounded sequence $(u_k) \in W^{1,1}(\Omega; \mathbb{R}^M)$. In other words, GYM is also the class of all generalized $W^{1,1}$ -gradient Young measures.

For generalized gradient Young measures satisfying $\lambda(\partial\Omega) = 0$, an explicit characterization is provided by the following result.

THEOREM 2.20 ([20, Theorem 9]). *Let $\Omega \subset \mathbb{R}^N$ be a bounded Lipschitz domain, and let $\Lambda := (\nu, \lambda, \nu^\infty) \in \mathcal{Y}_{\text{gen}}^1(\bar{\Omega}; \mathbb{R}^{M \times N})$ such that $\lambda(\partial\Omega) = 0$. Then $\Lambda \in \text{GYM}$ if and only if the following conditions hold:*

$$(i) \quad \int_{\Omega} \langle \nu_x, |\cdot| \rangle dx + \int_{\Omega} \langle \nu_x^\infty, |\cdot| \rangle d\lambda(x) < \infty,$$

and there exists $u \in \text{BV}(\Omega; \mathbb{R}^M)$ and $\omega_i \subset \Omega$, $\mathcal{L}^N(\omega_i) = 0$, such that for all $v : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ continuous and quasiconvex with at most linear growth,

$$(ii) \quad v(\nabla u(x)) \leq \langle \nu_x, v \rangle + \langle \nu_x^\infty, v^\sharp \rangle \frac{d\lambda}{d\mathcal{L}^N}(x) \quad \text{for all } x \in \Omega \setminus \omega_i,$$

$$(iii) \quad v^\sharp \left(\frac{dDu^s}{d|Du^s|}(x) \right) |Du^s| \leq \langle \nu_x^\infty, v^\sharp \rangle \lambda^s|_{\Omega} \quad \text{as measures.}$$

Here λ^s denotes the singular part of λ with respect to the Lebesgue measure \mathcal{L}^N , $\frac{d\lambda}{d\mathcal{L}^N}$ is the density of the absolutely continuous part of λ w.r.t. \mathcal{L}^N , and v^\sharp is the generalized recession function of v as introduced in (2.6).

Remark 2.21. As already pointed out in [20], (ii) and (iii) imply that Du is the center of mass of Λ restricted to Ω , i.e., $dDu(x) = \langle \nu_x, id \rangle dx + \langle \nu_x^\infty, id \rangle d\lambda(x)$ on Ω .

More recently, it was shown in [18, 19] that some of the conditions appearing in Theorem 2.20 can be simplified, and a smaller class of test functions v can be used, namely,

$$(2.23) \quad \text{SQ} := \left\{ v : \mathbb{R}^{M \times N} \rightarrow \mathbb{R} \mid \begin{array}{l} v \text{ is Lipschitz, quasiconvex, } v(A) \geq |A| \text{ for all } A \in \mathbb{R}^{M \times N}, \\ \text{and there exists } r_v > 0 \text{ s.t. } v(A) = v^\sharp(A) \text{ for all } |A| \geq r_v \end{array} \right\}.$$

Notice that functions $v \in \text{SQ}$ always have a recession function in the sense of (2.5), and $v^\sharp = v^\infty$. The crucial new observation proven in [18, 19] is that quasiconvex, positively one-homogeneous functions (like v^\sharp above) are automatically convex at zero and at all matrices of rank 1. In particular, this means that (iii) can essentially be dropped:

THEOREM 2.22 ([19, Theorem 6.2]). *Let $\Omega \subset \mathbb{R}^N$ be a bounded Lipschitz domain, and let $\Lambda := (\nu, \lambda, \nu^\infty) \in \mathcal{Y}_{\text{gen}}^1(\bar{\Omega}; \mathbb{R}^{M \times N})$ such that $\lambda(\partial\Omega) = 0$. Then $\Lambda \in \text{GYM}$ if*

and only if there exists $u \in \text{BV}(\Omega; \mathbb{R}^M)$ such that the following conditions hold:

- (0) Du is the center of mass of Λ on Ω ,
- (i) $\int_{\Omega} \langle \nu_x, |\cdot| \rangle dx + \int_{\bar{\Omega}} \langle \nu_x^\infty, |\cdot| \rangle d\lambda(x) < \infty$,

and there exists $\omega_i \subset \Omega$, $\mathcal{L}^N(\omega_i) = 0$, such that for all $v \in \text{SQ}$,

- (ii) $v(\nabla u(x)) \leq \langle \nu_x, v \rangle + \langle \nu_x^\infty, v^\infty \rangle \frac{d\lambda}{d\mathcal{L}^N}(x)$ for all $x \in \Omega \setminus \omega_i$.

Remark 2.23. Condition (i) is strictly speaking part of our definition of $\mathcal{Y}_{\text{gen}}^1$ (see subsection 2.6.2), and it is also part of the definition of \mathbf{Y} in the notation of [19]. In our case, it follows from the requirement that each element of $\mathcal{Y}_{\text{gen}}^1$ is generated by a bounded sequence in L^1 by using $v = |\cdot| = v^\infty$ and $g = 1$ in (2.13). Therefore, (i) could actually be dropped in Theorem 2.20 and Theorem 2.22, as well as in our main result, Theorem 3.1 below. But since it is not explicitly stated in our definition of $\mathcal{Y}_{\text{gen}}^1$, we prefer to keep it in the theorems.

Remark 2.24. Unlike in Theorem 2.20, (0) appears explicitly in Theorem 2.22, which is necessary because without (iii) it no longer holds automatically. This also means that despite the absence of (iii), a generalized gradient Young measure Λ is not entirely free on sets of Lebesgue measure zero. In particular, in order to be a BV derivative as required in (0), the center of mass must vanish on all sets of Hausdorff dimension less than $N - 1$ (see [2], e.g.).

Remark 2.25. For the prototypical example of a sequence of gradients concentrating at a single point at the boundary as introduced in Example 2.13, i.e., (∇u_k) with $u_k(x) := k^{N-1}u(kx)$, $u \in W_0^{1,1}(B; \mathbb{R}^M)$ (here, we assume that $0 \in \partial\Omega$), the center of mass of the generalized Young measure generated by this sequence charges the single point $0 \in \partial\Omega$ whenever $\int_{D_\varrho} \nabla u dx \neq 0$. The latter is even generic since the trace of u is free on the flat part of the boundary of the half-ball D_ϱ . But in view of Remark 2.24, whenever a generalized Young measure $(\nu_x, \lambda, \nu_x^\infty)$ satisfies $\lambda(\{x_0\}) > 0$ (with some $x_0 \in \partial\Omega$, $x_0 = 0$ in the example) and $0 \neq \int_{S^{MN-1}} s d\nu_{x_0}^\infty(s)$ ($= \|\nabla u\|_{L^1(D_\varrho; \mathbb{R}^M \times \mathbb{R}^N)}^{-1} \int_{D_\varrho} \nabla u dx$ in the example), it cannot be extended to a generalized gradient Young measure on a larger domain containing x_0 in its interior, because its center of mass charges the singleton $\{x_0\}$. Hence, in general, a generalized Young measures generated by a sequence of gradients charging $\partial\Omega$ (which corresponds to $\lambda(\partial\Omega) > 0$) cannot be extended to a generalized gradient Young measure on a larger domain.

3. The main result. We prove the following theorem which fully characterizes elements of GYM.

THEOREM 3.1 (Characterization of GYM up to the boundary). *Let $\Omega \subset \mathbb{R}^N$ be a bounded domain of class C^1 , and let $\Lambda := (\nu, \lambda, \nu^\infty) \in \mathcal{Y}_{\text{gen}}^1(\Omega; \mathbb{R}^{M \times N})$. Then $\Lambda \in \text{GYM}$ if and only if there exists $u \in \text{BV}(\Omega; \mathbb{R}^M)$ such that the following conditions hold:*

- (0) Du is the center of mass of Λ on Ω , i.e.,
 $dDu(x) = \langle \nu_x, id \rangle dx + \langle \nu_x^\infty, id \rangle d\lambda(x)$ on Ω ,
- (i) $\int_{\Omega} \langle \nu_x, |\cdot| \rangle dx + \int_{\bar{\Omega}} \langle \nu_x^\infty, |\cdot| \rangle d\lambda(x) < \infty$,

there exists $\omega_i \subset \Omega$, $\mathcal{L}^N(\omega_i) = 0$, such that for all $v \in \text{SQ}$ (cf. (2.23)),

$$(ii) \quad v(\nabla u(x)) \leq \langle \nu_x, v \rangle + \langle \nu_x^\infty, v^\infty \rangle \frac{d\lambda}{d\mathcal{L}^N}(x) \quad \text{for all } x \in \Omega \setminus \omega_i,$$

and there is $\omega_b \subset \partial\Omega$, $\lambda(\omega_b) = 0$ such that

$$(iii) \quad 0 \leq \langle \nu_x^\infty, v^\infty \rangle \quad \text{for all } x \in \partial\Omega \setminus \omega_b \text{ and all } v \in \Upsilon \cap \text{QSLB}(\varrho(x)),$$

where $\varrho(x)$ is the outer unit normal to $\partial\Omega$ at x .

Remark 3.2. If $N = 1$, it is clear that $\text{GYM} = \mathcal{Y}_{\text{gen}}^1$, simply because all functions on a one-dimensional domain are gradients of a potential. In Theorem 3.1, (0) then just determines u up to a constant, and given (0), (i)–(iii) always hold for $N = 1$. More precisely, (i) and (ii) then are a consequence of Jensen’s inequality, since quasi-convexity reduces to convexity. Similarly, (iii) becomes trivial because $v \in \text{QSLB}$ is equivalent to $v^\infty \geq 0$ for $N = 1$.

By Remark 3.2, it suffices to prove Theorem 3.1 for $N \geq 2$. The proof starts in section 4 below, where we split any given $\Lambda \in \mathcal{Y}_{\text{gen}}^1$ into two essentially disjoint pieces: an inner part Λ_i and a boundary part Λ_b . Due to Proposition 4.2, it will be enough to find conditions characterizing $\Lambda_i \in \text{GYM}$ and $\Lambda_b \in \text{GYM}$, separately. By Theorem 2.22 and the definition of Λ_i , conditions (0)–(ii) characterize Λ_i as an element of GYM. The proof of Theorem 3.1 will be completed by showing that (iii) is a necessary and sufficient condition for $\Lambda_b \in \text{GYM}$ (see section 5 and section 6).

4. Separating boundary and interior. The aim of this section is to show that when studying generalized gradient Young measures, it is enough to look at interior and boundary parts separately. For this purpose, we decompose $\Lambda := (\nu, \lambda, \nu^\infty) \in \mathcal{Y}_{\text{gen}}^1$ into two parts: the inner part $\Lambda_i = (\nu_i, \lambda_i, \nu_i^\infty)$ and the boundary part $\Lambda_b = (\nu_b, \lambda_b, \nu_b^\infty)$ defined as follows:

$$(4.1) \quad \langle \langle \Lambda_i, g \otimes v \rangle \rangle := \int_{\Omega} g(x) \langle \nu_x, v \rangle dx + \int_{\Omega} g(x) \langle \nu_x^\infty, v^\infty \rangle d\lambda(x) \quad \text{for all } g \in C(\bar{\Omega}), v \in \Upsilon,$$

i.e., $\Lambda|_{\Omega} = \Lambda_i|_{\Omega}$ and $\lambda_i(\partial\Omega) = 0$, and

$$(4.2) \quad \langle \langle \Lambda_b, g \otimes v \rangle \rangle := \int_{\partial\Omega} g(x)v(0) dx + \int_{\partial\Omega} g(x) \langle \nu_x^\infty, v^\infty \rangle d\lambda(x) \quad \text{for all } g \in C(\bar{\Omega}), v \in \Upsilon,$$

i.e., $\Lambda|_{\partial\Omega} = \Lambda_b|_{\partial\Omega}$ while $\Lambda_b|_{\Omega}$ is the “trivial” measure $(\delta_0, 0, -)$ generated by the constant sequence $0 \in \mathbb{R}^{M \times N}$ (here, recall that ν_x^∞ can be ignored on sets with λ -measure zero). Notice that (cf. (2.15))

$$(4.3) \quad \langle \langle \Lambda, g(x)v \rangle \rangle = \langle \langle \Lambda_i, g(x)v \rangle \rangle + \langle \langle \Lambda_b, g(x)v \rangle \rangle - \int_{\Omega} g(x)v(0) dx.$$

Remark 4.1. In a sense, the term $\int_{\Omega} g(x)v(0) dx$ in the definition (4.2) of $\Lambda_b = (\nu_b, \lambda_b, \nu_b^\infty)$ is artificial. Without it, however, $\nu_{b,x}$ would be the zero measure, and to have $\Lambda_b \in \mathcal{Y}_{\text{gen}}^1$, we need in particular $\nu_{b,x}$ to be a probability measure for a.e. $x \in \Omega$. We therefore choose $\nu_{b,x} = \delta_0$ for $x \in \Omega$, which corresponds to a generating sequence purely concentrating at the boundary.

PROPOSITION 4.2. *Let $\Lambda := (\nu, \lambda, \nu^\infty) \in \mathcal{Y}_{\text{gen}}^1$. Then $\Lambda \in \text{GYM}$ if and only if both $\Lambda_i \in \text{GYM}$ and $\Lambda_b \in \text{GYM}$.*

Proof. “only if”: Assuming that $\Lambda \in \text{GYM}$, let us see that $\Lambda_i \in \text{GYM}$ and $\Lambda_b \in \text{GYM}$. For this purpose let $(u_n) \subset \text{BV}(\Omega; \mathbb{R}^M)$ be a bounded sequence such that (Du_n) generates Λ . Without loss of generality, we may assume that $u_n \rightharpoonup^* u$ with some $u \in \text{BV}$. Using Lemma 2.2 with $J = 2$, $K_1 := \partial\Omega$, and $K_2 := \bar{\Omega}$, we decompose $\tilde{u}_n := u_n - u$ (or a suitable subsequence, not relabeled) as $\tilde{u}_n = c_n + d_n$ ($c_n = u_{1,n}$ and $d_n = u_{2,n}$ in Lemma 2.2).

We claim that (Dc_n) and $(Du + Dd_n)$ generate Λ_b and Λ_i , respectively. To prove the claim let $\Lambda_1 = (\nu_{1,x}, \lambda_1, \nu_{1,x}^\infty) \in \mathcal{Y}_{\text{gen}}^1$ be generated by (a subsequence not relabeled) of (Dc_n) , and analogously let $\Lambda_2 = (\nu_{2,x}, \lambda_2, \nu_{2,x}^\infty) \in \mathcal{Y}_{\text{gen}}^1$ be generated by (a subsequence not relabeled) of $(Du + Dd_n)$. Since all subsequences of (Dc_n) and $(Du + Dd_n)$ have another subsequence generating some generalized Young measure, it suffices to show that

$$\Lambda_1 = \Lambda_b \text{ and } \Lambda_2 = \Lambda_i.$$

We start by observing that as a consequence of (4.3), the fact that (Du_n) generates Λ , Proposition 2.3, and Proposition 2.4, then for every $g \in C(\bar{\Omega})$ and every $v \in \Upsilon$, we have that

$$\begin{aligned} (4.4) \quad & \langle\langle \Lambda_i, g \otimes v \rangle\rangle + \langle\langle \Lambda_b, g \otimes v \rangle\rangle - \int_{\Omega} g(x)v(0) \, dx \\ &= \langle\langle \Lambda, g \otimes v \rangle\rangle = \lim_n \int_{\Omega} g(x)dv(Du + Dd_n + Dc_n)(x) \\ &= \lim_n \int_{\Omega} g(x)dv(Du + Dd_n)(x) + \lim_n \int_{\Omega} g(x)dv(Du + Dc_n)(x) - \int_{\Omega} g(x)dv(Du)(x) \\ &= \lim_n \int_{\Omega} g(x)dv(Du + Dd_n)(x) + \lim_n \int_{\Omega} g(x)dv(Dc_n)(x) - \int_{\Omega} g(x)v(0) \, dx \\ &= \langle\langle \Lambda_2, g(x)v \rangle\rangle + \langle\langle \Lambda_1, g(x)v \rangle\rangle - \int_{\Omega} g(x)v(0) \, dx. \end{aligned}$$

Moreover, since $\{c_n \neq 0\} \subset (\partial\Omega)_{\frac{1}{n}}$, it is clear that for every $v \in \Upsilon$ and every $g \in C_c(\Omega)$,

$$\langle\langle \Lambda_1, g(x)v \rangle\rangle = \lim_n \int_{\Omega} g(x)dv(Dc_n)(x) = \int_{\Omega} g(x)v(0) \, dx,$$

whence $\nu_{1,x} = \nu_{b,x} = \delta_0$ for a.e. $x \in \Omega$ and $\lambda_1|_{\Omega} = \lambda_b|_{\Omega} = 0$, meaning that $\Lambda_1|_{\Omega} = (\delta_0, 0, 0) = \Lambda_b|_{\Omega}$. In view of (4.4) with $g \in C_c(\Omega)$, this also implies that $\Lambda_2|_{\Omega} = \Lambda_i|_{\Omega} = \Lambda|_{\Omega}$.

On the other hand, take now $g \in C(\partial\Omega)$ and extend it (without changing its name) to a function $g \in C(\bar{\Omega})$ by the Tietze theorem, and choose a sequence $(g_k)_{k \in \mathbb{N}} \subset C(\bar{\Omega})$ such that $g_k = g$ on $\partial\Omega$, $|g_k| \leq |g|$ on Ω , and $\text{supp } g_k \subset (\partial\Omega)_{\frac{1}{k}}$. Since Λ_2 is generated by $(Du + Dd_n)$, we have for every k

$$\begin{aligned} |\langle\langle \Lambda_2, g_k(x) \cdot |\cdot| \rangle\rangle| &= \left| \lim_n \int_{\Omega} g_k(x) \, d|Du + Dd_n|(x) \right| \\ &\leq C \|g_k\|_{C(\Omega)} \sup_n \int_{\text{supp } g_k} (d|Du|(x) + d|Dd_n|(x)) \xrightarrow[k \rightarrow +\infty]{} 0 \end{aligned}$$

by dominated convergence and the fact that (Dd_n) does not charge $\partial\Omega$. Again by dominated convergence, we infer that

$$\int_{\partial\Omega} g(x)d\lambda_2(x) = \lim_k \langle \langle \Lambda_2, g_k(x) | \cdot \rangle \rangle = 0.$$

Since this holds for arbitrary $g \in C(\partial\Omega)$, we get $\lambda_2(\partial\Omega) = 0 = \lambda_i(\partial\Omega)$, meaning that $\Lambda_2|_{\partial\Omega} = \Lambda_i|_{\partial\Omega}$. In view of (4.4), we conclude that $\Lambda|_{\partial\Omega} = \Lambda_1|_{\partial\Omega}$.

“if”: Assuming that $\Lambda_i \in \text{GYM}$ and $\Lambda_b \in \text{GYM}$, let us see that $\Lambda \in \text{GYM}$. Consider for that $(u_n^{(i)}), (u_n^{(b)}) \subset W^{1,1}(\Omega; \mathbb{R}^M)$ two sequences such that $(\nabla u_n^{(i)})$ generates Λ_i and $(\nabla u_n^{(b)})$ generates Λ_b . Choosing subsequences if necessary, we may assume that $u_n^{(i)} \xrightarrow{*} u^{(i)}$ and $u_n^{(b)} \xrightarrow{*} u^{(b)}$ in BV. Moreover, since $\nu_{b,x} = \delta_0$ for a.e. $x \in \Omega$ and $\lambda_b(\Omega) = 0$, it is clear that $Du^{(b)} = 0$ in Ω . Removing a constant if necessary, which does not affect the measure generated by the gradients, we can modify $u_n^{(b)}$ so that $u^{(b)} = 0$. We can decompose both sequences (or suitable subsequences) according to Lemma 2.2 (again with $J = 2$, $K_1 = \partial\Omega$, and $K_2 = \bar{\Omega}$) into a part c_n purely concentrating at the boundary and an interior remainder d_n which does not charge $\partial\Omega$:

$$u_n^{(i)} - u^{(i)} = c_n^{(i)} + d_n^{(i)} \quad \text{and} \quad u_n^{(b)} = c_n^{(b)} + d_n^{(b)}.$$

Similarly to the first part of the proof, $(Dd_n^{(i)} + Du^{(i)})$ still generates $(\Lambda_i)_i = \Lambda_i$, and $(Dc_n^{(b)})$ still generates $(\Lambda_b)_b = \Lambda_b$. Moreover, Proposition 2.3 applies to the decomposition

$$w_n := c_n^{(b)} + d_n^{(i)},$$

and together with Proposition 2.4 we get

$$\begin{aligned} & \lim_n \int_{\Omega} g(x)dv(Du^{(i)} + Dw_n) \\ &= \lim_n \left(\int_{\Omega} g(x)dv(Du^{(i)} + Dd_n^{(i)}) + \int_{\Omega} g(x)dv(Dc_n^{(b)}) - \int_{\Omega} g(x)v(0) dx \right) \\ &= \langle \langle \Lambda_i, g(x)v \rangle \rangle + \langle \langle \Lambda_b, g(x)v \rangle \rangle - \int_{\Omega} g(x)v(0) dx = \langle \langle \Lambda, g \otimes v \rangle \rangle \end{aligned}$$

in view of (4.3), which proves that $(Du^{(i)} + Dw_n)$ generates Λ . □

5. Necessary conditions at the boundary. In view of Proposition 4.2 (see also the end of section 3) we will restrict ourselves to investigating $\Lambda_b = (\delta_0, \lambda(\partial\Omega \cap \cdot), \nu_x^\infty) \in \mathcal{Y}_{\text{gen}}^1$ (cf. (4.2)). Necessary conditions for Λ_b to be a generalized gradient Young measure will be derived with the help of Proposition 2.8 and Remark 2.19. In fact, on a planar piece of the boundary with a fixed outer normal ϱ , the next proposition is a straightforward consequence of Proposition 2.8, and using separability of $\text{QSLB}(\varrho)$ (as a subset of the continuous, positively one-homogeneous functions with the topology of $C(S^{MN-1})$, cf. Remark 2.6) it is easy to see that the set ω_b (in condition (iii) of Theorem 3.1) can be chosen independently of the test functions $v \in \text{QSLB}(\varrho)$. However, once the boundary is not part of a hyperplane, the set of test functions $\text{QSLB}(\varrho(x))$ depends on $x \in \partial\Omega$, which significantly complicates the definition of ω_b .

PROPOSITION 5.1. *If $\Lambda^b \in \text{GYM}$ then there is $\omega_b \subset \partial\Omega$, $\lambda(\omega_b) = 0$ such that*

$$0 \leq \langle \nu_x^\infty, v^\infty \rangle$$

for all $x \in \partial\Omega \setminus \omega_b$ and all $v \in \Upsilon \cap \text{QSLB}(\varrho(x))$.

Proof. Suppose that $(\nabla c_n)_n$ generates Λ_b for a bounded sequence $(c_n) \subset W^{1,1}(\Omega; \mathbb{R}^M)$, which in particular implies that $c_n \overset{*}{\rightharpoonup} 0$ in $BV(\Omega; \mathbb{R}^M)$.

Given $j \in \mathbb{N}$, let us consider $x_i^j \in \partial\Omega$, $i = 1, \dots, I_j$, such that

$$\partial\Omega \subset \bigcup_{i=1}^{I_j} B(x_i^j, 1/j).$$

Fix $j \in \mathbb{N}$ and $i \in \{1, \dots, I_j\}$, and suppose there is $v \in \Upsilon$ such that $v \in \text{QSLB}(y)$ for all $y \in \partial\Omega \cap B(x_i^j, 1/j)$ (as we will see below, such v exist at least for large j , essentially due to Lemma 2.9(ii)). For every such v , we have the following: Take $g \in C_0(B(x_i^j, 1/j))$, $0 \leq g$, and extend it by zero to the whole $\bar{\Omega}$. Proposition 2.8 applied to $f : (x, A) \mapsto g(x)v(A)$ asserts that

$$(5.1) \quad \lim_{n \rightarrow \infty} \int_{\Omega} g(x)v(\nabla c_n(x)) \, dx \geq \int_{\Omega} g(x)v(0) \, dx.$$

On the other hand, since $(\nabla c_n)_n$ generates $\Lambda_b = (\delta_0, \lambda(\partial\Omega \cap \cdot), \nu_x^\infty)$,

$$\lim_{n \rightarrow \infty} \int_{\Omega} g(x)v(\nabla c_n) \, dx = \int_{\Omega} g(x)v(0) \, dx + \int_{\bar{\Omega}} \int_{S^{M \times N-1}} g(x)v^\infty(A) d\nu_x^{b,\infty}(A) d\lambda(x),$$

which from (5.1) implies that

$$(5.2) \quad \int_{\bar{\Omega}} g(x)\langle \nu_x^{b,\infty}, v^\infty \rangle d\lambda(x) = \int_{\bar{\Omega} \cap B(x_i^j, 1/j)} g(x)\langle \nu_x^{b,\infty}, v^\infty \rangle d\lambda(x) \geq 0.$$

As $g \in C_0(B(x_i^j, 1/j))$, $0 \leq g$, is arbitrary, we conclude that

$$\langle \nu_x^{b,\infty}, v^\infty \rangle \geq 0$$

for $x \in \partial\Omega \cap B(x_i^j, 1/j) \setminus E_{i,j,v}$ for some set $E_{i,j,v} \subset \partial\Omega \cap B(x_i^j, 1/j)$ with $\lambda(E_{i,j,v}) = 0$.

Let $\{v_k\}_{k \in \mathbb{N}}$ be a dense subset of $C(S^{M \times N-1})$. Given $\varepsilon > 0$ define

$$E_{i,j,\varepsilon} := \bigcup \left\{ E_{i,j,v_k(\cdot) + \varepsilon|\cdot|} : k \in \mathbb{N} \text{ such that } v_k(\cdot) + \varepsilon|\cdot| \in \text{QSLB}(y) \text{ for all } y \in \partial\Omega \cap B(x_i^j, 1/j) \right\}$$

and observe that $\lambda(E_{i,j,\varepsilon}) = 0$. For each $m \in \mathbb{N}$ consider now $\varepsilon_m = 1/m$ and define $E = \bigcup_{i,j,m} E_{i,j,m}$. Let us see that the claim holds taking $\omega^b = E$.

For this purpose we choose a sequence $0 < \varepsilon_m \rightarrow 0$. Moreover, let $x_0 \in \partial\Omega \setminus \omega^b$ and $v \in \Upsilon \cap \text{QSLB}(x_0)$. Given m , by Lemma 2.9(ii), there exists j large enough such that $v(\cdot) + \frac{\varepsilon_m}{2}|\cdot| \in \text{QSLB}(y)$ for all $y \in B(x_i^j, 1/j)$, where i is chosen such that $x_0 \in B(x_i^j, 1/j)$. Let now $\{v_{k_l}\}$ be a subsequence of $\{v_k\}$ such that $|v_{k_l} - v| < \frac{\varepsilon_m}{2}$ in $C(S^{M \times N-1})$. Then $v_{k_l}(\cdot) + \varepsilon_m|\cdot| \in \text{QSLB}(y)$ for all $y \in B(x_i^j, 1/j)$ and thus

$$\langle \nu_{x_0}^{b,\infty}, v_{k_l}(\cdot) + \varepsilon_m|\cdot| \rangle \geq 0.$$

The claim follows by letting $l, m \rightarrow +\infty$. □

6. Sufficient conditions at the boundary. The objective of this section is to prove that condition (iii) in Theorem 3.1 allows for a construction of a sequence of gradients generating $\Lambda_b = (\delta_0, \lambda(\partial\Omega \cap \cdot), \nu_x^\infty) \in \mathcal{Y}_{\text{gen}}^1$ (cf. (4.2)). More precisely we will derive the following result.

PROPOSITION 6.1. *Assume that for λ -almost all $x \in \partial\Omega$ we have that*

$$0 \leq \int_{S^{M \times N-1}} v^\infty(A) \nu_x^\infty(dA)$$

for all $v \in \Upsilon \cap \text{QSLB}(\varrho(x))$. Then $\Lambda_b \in \text{GYM}$.

To prove Proposition 6.1 (at the end of this section) we follow ideas developed in [23] and [13], relying on the Hahn–Banach theorem, allowing us to show that all GYM-measures that concentrate at the boundary can be approximated in the weak* convergence by a sequence of “elementary” measures.

Below, we often work with functions $v \in C(S^{M \times N-1})$, which we extend to the whole space as a positively one-homogeneous function (without changing its name) so that in particular $v = v^\infty$. Given a unit vector ϱ in \mathbb{R}^N , we start by defining two sets of measures:

$$A^\varrho := \{ \mu \in \mathcal{M}(S^{M \times N-1}); \mu \geq 0, \langle \mu, v \rangle \geq 0 \text{ for all } v \in C(S^{M \times N-1}) \\ \text{s.t. } v^\infty \in \text{QSLB}(\varrho) \}$$

and

$$H^\varrho := \{ \bar{\delta}_{\varrho, \nabla u}, \quad u \in W_0^{1,1}(B; \mathbb{R}^M) \},$$

where for all $v \in C(S^{M \times N-1})$, given $u \in W_0^{1,1}(B; \mathbb{R}^M)$, the measure $\bar{\delta}_{\varrho, \nabla u}$ is defined by

$$\langle \bar{\delta}_{\varrho, \nabla u}, v \rangle := \int_{D_\varrho} v \left(\frac{\nabla u(x)}{|\nabla u(x)|} \right) |\nabla u(x)| \, dx.$$

Both H^ϱ and A^ϱ are sets of measures on the unit sphere in $\mathbb{R}^{M \times N}$ and, in addition, we observe that

$$(6.1) \quad H^\varrho \subset A^\varrho.$$

Indeed to see (6.1) let $\bar{\delta}_{\varrho, \nabla u} \in H^\varrho$ with $u \in W_0^{1,1}(B; \mathbb{R}^M)$. Hence

$$\langle \bar{\delta}_{\varrho, \nabla u}, v \rangle = \int_{D_\varrho} v^\infty(\nabla u(x)) \, dx,$$

and if $v \in \text{QSLB}(\varrho)$ then (see 2.9) it holds that $\int_{D_\varrho} v^\infty(\nabla u(x)) \, dx \geq 0$ for all $u \in W_0^{1,1}(B; \mathbb{R}^M)$, i.e., $\bar{\delta}_{\varrho, \nabla u} \in A^\varrho$.

LEMMA 6.2. *Let $N \geq 2$. Then the set H^ϱ is convex.*

Proof. Consider $u_1, u_2 \in W_0^{1,1}(B; \mathbb{R}^M)$. We need to show that for any $0 \leq t \leq 1$ the convex combination $t\bar{\delta}_{\varrho, \nabla u_1} + (1-t)\bar{\delta}_{\varrho, \nabla u_2} \in H^\varrho$. Take $x_0 \in B \cap \{x \in \mathbb{R}^N; \varrho \cdot x = 0\}$ such that $|x_0| = \frac{2}{3}$ and define

$$\tilde{u}_1(x) := 3^{N-1}u_1(3x) \text{ and } \tilde{u}_2(x) := 3^{N-1}u_2(3(x-x_0)).$$

Observe that $\tilde{u}_1 \in W_0^{1,1}(B(0, 1/3); \mathbb{R}^M)$, $\tilde{u}_2 \in W_0^{1,1}(B(x_0, 1/3); \mathbb{R}^M)$ and then extend these functions by zero to the whole \mathbb{R}^N (without changing its name), so that, in particular, $\tilde{u}_1, \tilde{u}_2 \in W_0^{1,1}(B; \mathbb{R}^M)$ and have disjoint supports. Let

$$u := t\tilde{u}_1 + (1-t)\tilde{u}_2(x).$$

Given $v \in C(S^{M \times N-1})$ extend it, as before, to the whole space as a positively one-homogeneous function. Thus

$$\begin{aligned} \int_{D_e} v(\nabla u(x)) \, dx &= t \int_{D_e \cap B(0,1/3)} v(3^N \nabla u_1(3x)) \, dx \\ &\quad + (1-t) \int_{D_e \cap B(x_0,1/3)} v(3^N \nabla u_1(3(x-x_0))) \, dx \\ &= t \int_{D_e} v(\nabla u_1(y)) \, dy + (1-t) \int_{D_e} v(\nabla u_2(y)) \, dy, \end{aligned}$$

which proves the claim. □

The next result is a standard application of the Hahn–Banach theorem.

PROPOSITION 6.3. *The set A^e is the weak* closure of H^e .*

Remark 6.4. The topological weak* closure and the sequential weak* closure of H^e coincide. The reason is that H^e is a subset of the positive cone $P := \{\mu \in \mathcal{M}(S^{MN-1}) \mid \mu \geq 0\}$ in $\mathcal{M}(S^{MN-1})$. For such sets, topological and sequential weak* closure always are the same, even if they are unbounded. To see that, take any $H \subset P$ and let $\mu \in A := \overline{H}^{w*}$, the topological weak* closure of H in $\mathcal{M}(S^{MN-1})$. It suffices to show that $\mu \in \overline{H}^{s-w*}$, the sequential weak* closure of H . There exists $t > 0$ such that

$$\mu \in N_t := \{\gamma \in \mathcal{M}(S^{MN-1}) \mid \int_{S^{MN-1}} 1 \, d\gamma = \langle \gamma, 1 \rangle < t\},$$

which by definition is a weak*-open set in $\mathcal{M}(S^{MN-1})$. As a consequence of the latter,

$$\mu \in N_t \cap \overline{H}^{w*} \subset \overline{N_t \cap H}^{w*}.$$

Moreover, $N_t \cap H$ is bounded in the norm of \mathcal{M} by the definition of N_t , because $H \subset P$ and, on P , $\langle 1, \cdot \rangle = \|\cdot\|_{\mathcal{M}}$. Since $\mathcal{M}(S^{MN-1})$ is the dual of the separable Banach space $C(S^{MN-1})$, the weak* topology is metrizable on bounded subsets of $\mathcal{M}(S^{MN-1})$, and therefore the sequential weak* closure and the topological weak* closure of $N_t \cap H$ are the same. We conclude that $\mu \in \overline{N_t \cap H}^{s-w*} \subset \overline{H}^{s-w*}$.

Proof of Proposition 6.3. Assume that $A^e \neq \overline{H^e}^{w*}$. As $H^e \subset A^e$ and A^e is clearly closed with respect to the weak* topology, this implies that there exists $\mu_0 \in A^e$ such that $\mu_0 \notin \overline{H^e}^{w*}$. By the Hahn–Banach theorem, we can separate μ_0 from the convex set H^e by a hyperplane, i.e., there exist $v \in C(S^{MN-1})$ and $a \in \mathbb{R}$ such that on the one hand

$$\langle \bar{\mu}_0, v \rangle < a,$$

while on the other hand

$$\langle \bar{\delta}_{e, \nabla u}, v \rangle \geq a$$

for all $\bar{\delta}_{e, \nabla u} \in H^e$. Extending v , as before, to the whole space as a positively one-homogeneous function, and noting that $v = v^\infty$, we get that

$$\int_{D_e} v^\infty(\nabla u(x)) \, dx \geq a \quad \text{for every } u \in W_0^{1,1}(B; \mathbb{R}^M).$$

By positive one-homogeneity, this forces $a \leq 0$. Moreover, there cannot be an admissible u such that $\int_{D_e} v^\infty(\nabla u(x)) \, dx < 0$, because multiplying this u by arbitrarily large

constants immediately gives a contradiction (a might be negative, but it is finite). Hence,

$$\int_{D_\varrho} v^\infty(\nabla u(x)) \, dx \geq 0 \quad \text{for every } u \in W_0^{1,1}(B; \mathbb{R}^M).$$

As a consequence, $v^\infty \in \text{QSLB}(\varrho)$ by Remark 2.6. Thus if $\mu \in A^\varrho$, we get $a > \langle \mu_0, v \rangle \geq 0 \geq a$, a contradiction. \square

Remark 6.5.

(i) By definition, for any given $u \in W_0^{1,1}(B; \mathbb{R}^M)$, $\bar{\delta}_{\varrho, \nabla u} \in H^\varrho$ if and only if the generalized Young measure $\Lambda := (\nu_x, \lambda, \nu_x^\infty)$ on the half-ball D_ϱ given by

$$\nu_x := \delta_0, \quad \lambda := \|\nabla u\|_{L^1(D_\varrho; \mathbb{R}^{M \times N})} \delta_0, \quad \nu_0^\infty := \frac{1}{\|\nabla u\|_{L^1(D_\varrho; \mathbb{R}^{M \times N})}} \bar{\delta}_{\varrho, \nabla u}$$

is generated by (∇u_k) , with $u_k(x) := k^{N-1}u(kx)$. Notice that (u_k) is a rather specific sequence “purely concentrating” at $0 \in \partial D_\varrho$, supported in balls of radius $\frac{1}{k}$.

(ii) On the other hand, suppose that $\Lambda := (\delta_0, c\delta_0, \nu_0^\infty)$, with some constant $c \geq 0$, is a generalized Young measure on D_ϱ such that $\mu := c\nu_0^\infty \in A^\varrho$. Proposition 6.3 and Remark 6.4 then imply that Λ is generated by a sequence (∇w_j) which is also “purely concentrating” at $0 \in \partial D_\varrho$. This can be seen using a diagonalization argument: Since A^ϱ is the weak* closure of H^ϱ , we can find a sequence $(u^{(j)}) \in W_0^{1,1}(B; \mathbb{R}^M)$ such that $\bar{\delta}_{\varrho, \nabla u^{(j)}} \rightharpoonup^* \mu = c\nu_0^\infty \in A^\varrho$. Then we can get an appropriate choice for w_j by letting $w_j(x) := u_{k(j)}^{(j)} = k(j)^{N-1}u^{(j)}(k(j)x)$ with $k(j) \rightarrow \infty$ fast enough as $j \rightarrow \infty$. Let us also point out that since $\bar{\delta}_{\varrho, \nabla u^{(j)}} \rightharpoonup^* c\nu_0^\infty$, we automatically have $\int_{D_\varrho} |\nabla u^{(j)}| \, dx \rightarrow c$ because ν_0^∞ is a probability measure on S^{MN-1} . Consequently, $\|\nabla w_j\|_{L^1} \rightarrow c$ on D_ϱ .

PROPOSITION 6.6. *Assume that $J \geq 1$, $\Lambda := (\delta_0, \nu^\infty, \lambda)$ such that $\lambda = \sum_{i=1}^J a_i \delta_{x_i}$, where $x_i \in \partial\Omega$ and $a_i > 0$ for $1 \leq i \leq J$, and $\nu_{x_i}^\infty \in A^{\varrho(x_i)}$ for $1 \leq i \leq J$ with $\varrho(x_i) \in \mathbb{R}^N$ the unit outer normal to $\partial\Omega$ at x_i . Then $\Lambda \in \text{GYM}$.*

Proof. First notice that $x \mapsto \nu_x^\infty$ is defined λ -a.e. in $\bar{\Omega}$. Assume first that $\nu_{x_i}^\infty \in H^{\varrho(x_i)}$ for all $1 \leq i \leq J$. Therefore, there is $u_i \in W_0^{1,1}(B; \mathbb{R}^M)$ such that, for every $v^\infty \in C(\mathbb{R}^{M \times N})$ positively one-homogeneous,

$$\int_{D_{\varrho(x_i)}} v^\infty(\nabla u_i(y)) \, dy = \int_{S^{M \times N - 1}} v^\infty(A) \, d\nu_{x_i}^\infty(A)$$

by definition of $H^{\varrho(x_i)}$. Consequently,

$$\begin{aligned} (6.2) \quad & \lim_{j \rightarrow \infty} a_i \int_{\Omega \cap B(x_i, 1/j)} v^\infty(\nabla u_i(j(x - x_i))) \, dx \\ &= \lim_{j \rightarrow \infty} a_i \int_{j(-x_i + \Omega \cap B(x_i, 1/j))} v^\infty(\nabla u_i(y)) \, dy \\ &= a_i \int_{D_{\varrho(x_i)}} v^\infty(\nabla u_i(y)) \, dy = a_i \int_{S^{M \times N - 1}} v^\infty(A) \, d\nu_{x_i}^\infty(A). \end{aligned}$$

Consequently, $(\sum_{i=1}^J a_i j^{N-1} u_i(j(x - x_i)))_{j \in \mathbb{N}} \subset W^{1,1}(\Omega; \mathbb{R}^M)$ is a sequence such that the gradients $(\sum_{i=1}^J a_i j^N \nabla u_i(j(x - x_i)))_{j \in \mathbb{N}}$ generate Λ .

Assume now that $\nu_{x_i}^\infty \in A^{\varrho(x_i)}$. From Proposition 6.3 and Remark 6.4, for each fixed $1 \leq i \leq J$ there exists $(u_i^k)_{k \in \mathbb{N}} \subset W_0^{1,1}(B; \mathbb{R}^M)$ such that, for every $v^\infty \in C(S^{M \times N-1})$ positively one-homogeneous,

$$\begin{aligned}
 (6.3) \quad & \lim_{k \rightarrow \infty} \lim_{j \rightarrow \infty} a_i \int_{\Omega \cap B(x_i, 1/j)} v^\infty(\nabla u_i^k(j(x - x_i))) \, dx \\
 &= \lim_{k \rightarrow \infty} \lim_{j \rightarrow \infty} a_i \int_{j(\Omega \cap B(x_i, 1/j) - x_i)} v^\infty(\nabla u_i^k(y)) \, dy \\
 &= a_i \int_{S^{M \times N-1}} v^\infty(A) \, d\nu_{x_i}^\infty(A).
 \end{aligned}$$

As $C(S^{M \times N-1})$ is separable and $\|\nabla u_i^k\|_{L^1(\Omega; \mathbb{R}^{M \times N})} \leq C$ for all $k \in \mathbb{N}$, the proof is finished by a diagonalization argument. \square

The property of quasi-sublinear growth from below is inevitably connected with the boundary normal. The following proposition shows that suitable rotations allow us to “average” along the boundary.

PROPOSITION 6.7. *Let $x_0 \in \partial\Omega$. Moreover, let $\{R(x)\}_{x \in \partial\Omega} \subset \text{SO}(N)$ be a family of rotation matrices such that $x \mapsto R(x)$ is continuous and bounded on a set $\Gamma \subset \partial\Omega$ and, for each $x \in \Gamma$, $\varrho(x) = R(x)\varrho(x_0)$. Given a measurable set $E \subset \Gamma$ such that $\lambda(E) > 0$, we define $\mu_{x_0} = \mu_{x_0, E} \in \mathcal{M}(S^{M \times N-1})$ as the measure that satisfies*

$$\int_{S^{M \times N-1}} v(A) \, d\mu_{x_0}^\infty(A) := \frac{1}{\lambda(E)} \int_E \int_{S^{M \times N-1}} v(AR^\top(x)) \, d\nu_x^\infty(A) \, d\lambda(x)$$

for every $v \in C(S^{M \times N-1})$. If $\{\nu_x^\infty\}_{x \in E}$ satisfies for every $v \in \text{QSLB}(x)$

$$(6.4) \quad 0 \leq \int_{S^{M \times N-1}} v(A) \, d\nu_x^\infty(A),$$

then for every $v \in \text{QSLB}(x_0)$

$$(6.5) \quad 0 \leq \int_{S^{M \times N-1}} v^\infty(A) \, d\mu_{x_0}^\infty(A).$$

Proof. For every $v \in \Upsilon_S(\mathbb{R}^{M \times N})$ which is qslb at x_0 , $A \mapsto v(AR_x^{-1})$ is qslb at x by Lemma 2.9. Hence, (6.4) implies (6.5) by the definition of μ_{x_0} . \square

Proof of Proposition 6.1. For each $n \in \mathbb{N}$ cover \mathbb{R}^N with a family of pairwise disjoint cubes of side length 2^{-n} , translations of $Q_{n,0} := [0, 2^{-n})^N$, and let $Q_{n,j}$, $j \in J(n)$, be the collection of those cubes Q in the family that satisfy $\lambda(Q \cap \partial\Omega) > 0$. Moreover, for each n and each $j \in J(n)$ let $E_{n,j} := Q_{n,j} \cap \partial\Omega$, choose a point $x_{n,j} \in E_{n,j}$, and choose a family of rotations $(R_{n,j}(x))_{x \in E_{n,j}} \subset \mathbb{R}^{N \times N}$ such that $R_{n,j}(x_{n,j}) = \mathbb{I}$ and $\varrho(x) = R_{n,j}(x)\varrho(x_{n,j})$ for every $x \in E_{n,j}$, where $\varrho(x)$ denotes the outer normal at $x \in \partial\Omega$, $x \mapsto R_{n,j}(x)$ is continuous on $\overline{E}_{n,j}$ and

$$(6.6) \quad \sup_{j \in J(n)} \sup_{x \in E_{n,j}} |R_{n,j}(x)^{-1} - \mathbb{I}| \xrightarrow{n \rightarrow \infty} 0,$$

which is possible since $\partial\Omega$ is of class C^1 . We define for each $n \in \mathbb{N}$

$$\theta_n(x) := \sum_{j \in J(n)} \lambda(E_{n,j}) \delta_{x_{n,j}}(x),$$

and, for every $v \in C(S^{M \times N-1})$ and every $j \in J(n)$,

$$\begin{aligned} & \int_{S^{M \times N-1}} v(A) d\eta_{n,x}^\infty(A) \\ & := \frac{1}{\lambda(E_{n,j})} \int_{E_{n,j}} \int_{S^{M \times N-1}} v(A R^\top(y)) d\nu_y^\infty(A) d\lambda(y) \quad \text{if } x=x_{n,j}. \end{aligned}$$

Here, note that for $x \in \partial\Omega \setminus \{x_{n,j} \mid j \in J(n)\}$, the definition of $\eta_{n,x}^\infty$ does not matter since $\theta_n(\partial\Omega \setminus \{x_{n,j} \mid j \in J(n)\}) = 0$. Clearly, $(\delta_0, \theta_n, \eta_{n,x}^\infty) \in \text{GYM}$, by Proposition 6.6 and Proposition 6.7. Finally, observe that, by (6.6) and uniform continuity of continuous functions on compact sets, it holds for any $g \in C(\bar{\Omega})$ and $v \in C(S^{M \times N-1})$ that

$$\begin{aligned} & \int_{\partial\Omega} g(x) \int_{S^{M \times N-1}} v(A) d\eta_{n,x}^\infty(A) d\theta_n(x) \\ & = \sum_{j \in J(n)} g(x_{n,j}) \int_{E_{n,j}} \int_{S^{M \times N-1}} v(A R_{n,j}^\top(y)) d\nu_y^\infty(A) d\lambda(y) \\ & \rightarrow \int_{\partial\Omega} g(y) \int_{S^{M \times N-1}} v(A) d\nu_y^\infty(A) d\lambda(y) \quad \text{as } n \rightarrow +\infty. \end{aligned}$$

As $(\delta_0, \theta_n, \eta_{n,x}^\infty) \in \text{GYM}$ for any $n \in \mathbb{N}$, there is for every n a sequence $(u_j^n)_{j \in \mathbb{N}} \subset W^{1,1}(\Omega; \mathbb{R}^M)$ such that $(\nabla u_j^n)_{j \in \mathbb{N}}$ generates $(\delta_0, \theta_n, \eta_{n,x}^\infty)$. Due to the fact that $\lim_{j \rightarrow \infty} \|\nabla u_j^n\|_{L^1} = \theta_n(\bar{\Omega}) = \lambda(\bar{\Omega})$, there exists $C > 0$ such that $\|\nabla u_j^n\|_{L^1} < C$ for all j, n (if necessary, modify u_j^n for finitely many j for each n). Moreover, adding suitable constants to u_j^n and using Poincaré’s inequality gives us $\|u_j^n\|_{W^{1,1}} < C$ for all j, n . Finally, we use the separability of $C(\bar{\Omega})$ and $C(S^{M \times N-1})$ to extract a diagonal subsequence that generates Λ_b : Let $G \subset C(\bar{\Omega})$ and $V \subset C(S^{M \times N-1})$ be countable dense subsets. Since $(\nabla u_j^n)_{j \in \mathbb{N}}$ generates $(\delta_0, \theta_n, \eta_{n,x}^\infty)$ for each n , there exists a sequence $j(n) \rightarrow \infty$ such that

$$\int_{\Omega} g(x) v(\nabla u_{j(n)}^n) dx \rightarrow \langle \langle \Lambda_b, g \otimes v \rangle \rangle \quad \text{for every } g \in G, v \in V,$$

where v is understood to be extended to $\mathbb{R}^{M \times N}$ as a positively one-homogeneous function. Together with the fact that $(\nabla u_{j(n)}^n)$ is bounded in L^1 , we infer that $(\nabla u_{j(n)}^n)$ generates Λ_b , whence $\Lambda_b \in \text{GYM}$. Here, notice that since the Young measure part of Λ_b is δ_0 and $\lambda(\Omega) = 0$, it suffices to test the generating relation with positively one-homogeneous functions v , as using $v = |\cdot|$ (or anything close to that) suffices to check that the Young measure part really is δ_0 and the sequence really only charges ∂O . \square

Applying Theorem 2.22 to Λ_i defined in (4.1) we see that Λ from Theorem 3.1 can be written as $\Lambda = \Lambda_i + \Lambda_b$, where Λ_i and Λ_b are as in Proposition 4.2. Altogether, Theorem 3.1 is proved.

7. The Souček space W_μ^1 and its relation to BV and GYM. Nowadays, it is well known that the extension of a functional with linear growth in $W^{1,1}$ subject to Dirichlet boundary conditions leads to an unconstrained functional on BV, which, instead of the boundary condition, has an extra term at the boundary penalizing the difference of the “inner” trace and the “true” boundary value. Mostly forgotten by the modern literature covering this topic, J. Souček introduced and analyzed the larger

space W_μ^1 in [30]¹ with a similar application in mind. Since generalized BV-gradient Young measures are actually naturally embedded in W_μ^1 , we want to provide a short overview here. Besides, for the toy problem featured in the introduction, W_μ^1 is a more natural space than BV. As we shall see below, $W_\mu^1(\bar{\Omega})$ is actually isomorphic to $BV(\Omega) \times \mathcal{M}(\partial\Omega)$. Here, the second component is a measure representing an artificial “outer” trace on the boundary.

Just as in [30], we assume throughout this section that $\Omega \subset \mathbb{R}^N$ is a bounded domain of class C^1 .

DEFINITION 7.1 (Souček space W_μ^1 , [30, Definition 1]). *We define $W_\mu^1(\bar{\Omega})$ as the space of those $(u, \alpha) \in L^1(\Omega) \times \mathcal{M}(\bar{\Omega}; \mathbb{R}^M)$ which are in the weak* closure of $W^{1,1}(\Omega)$ on $\bar{\Omega}$ in the sense that for a suitable sequence $(u_n) \in W^{1,1}(\Omega)$,*

$$u_n \rightarrow u \text{ in } L^1(\Omega) \text{ and } \int_\Omega \psi \cdot \nabla u_n \, dx \rightarrow \int_{\bar{\Omega}} \psi \cdot d\alpha(x) \text{ for all } \psi \in C(\bar{\Omega}; \mathbb{R}^N).$$

The norm of W_μ^1 is given by

$$\|(u, \alpha)\|_{W_\mu^1(\bar{\Omega})} := \|u\|_{L^1(\Omega)} + \sum_{i=1}^N \|\alpha_i\|_{\mathcal{M}(\bar{\Omega})}, \text{ where } \alpha = (\alpha_1, \dots, \alpha_N).$$

From the modern perspective, it is clear that any such u is an element of $BV(\Omega)$, and if we restrict the vector-valued measure α to Ω , it coincides with the BV derivative Du . In that sense, W_μ^1 can be naturally projected onto BV. However, α carries extra information on $\partial\Omega$, and it turns out that $\alpha|_{\partial\Omega}$ is exactly the derivative of a jump at $\partial\Omega$ to some measure-valued “outer” trace that in particular captures the boundary value along sequences satisfying a fixed Dirichlet condition.

More precisely, Souček shows that any $(u, \alpha) \in W_\mu^1(\bar{\Omega})$ has two different traces on $\partial\Omega$: The first one is the *outer trace* $\beta \in \mathcal{M}(\partial\Omega)$ characterized by the version of Green’s formula derived in [30, Theorem 1] (with ϱ denoting the outer normal on $\partial\Omega$):

$$(7.1) \quad \int_{\partial\Omega} \varphi \varrho \, d\beta = \int_\Omega u \nabla \varphi \, dx + \int_{\bar{\Omega}} \varphi \, d\alpha \text{ for every } \varphi \in C^1(\bar{\Omega}).$$

The outer trace is weakly* continuous with respect to weak* convergence of (u, α) in $\mathcal{M}(\bar{\Omega}) \times \mathcal{M}(\bar{\Omega}; \mathbb{R}^N)$ [30, Theorem 2], in strong contrast to the situation in BV. Here, observe that the kind of behavior of a sequence that has to be ruled out in BV to ensure continuity of the trace (usually by imposing strict convergence); namely, a jump moving to or growing at the boundary is not lost in the weak* limit in $W_\mu^1(\Omega)$ as it appears in $\alpha|_{\partial\Omega}$. In addition, in [30, Definition 4] the *inner trace* $\beta^0 \in \mathcal{M}(\partial\Omega)$ of $(u, \alpha) \in W_\mu^1$ is defined as the outer trace of $(u, \bar{\alpha})$, where $\bar{\alpha}$ is obtained from α by dropping the contribution on $\partial\Omega$, i.e., $\bar{\alpha}|_\Omega := \alpha|_\Omega = Du$ and $\bar{\alpha}|_{\partial\Omega} := 0$. Here, $(u, \bar{\alpha}) \in W_\mu^1$ due to [30, Theorem 8] (alternatively, one can use the fact that the BV function u is the limit of a strictly convergent sequence $(u_n) \subset W^{1,1}$ and observe that strict convergence implies that $\nabla u_n \rightharpoonup^* \bar{\alpha}$ in $\mathcal{M}(\bar{\Omega})$, whence $(u, \bar{\alpha}) \in W_\mu^1$). Due to (7.1) and the fact that $\alpha = Du$ on Ω , β^0 coincides with trace of u in the sense of BV. As a consequence (or, alternatively, due to [30, Theorem 10]), the inner

¹Here, we stick to the original notation $W_\mu^1(\bar{\Omega})$ for the space of [30], although using $BV(\bar{\Omega})$ instead would also make sense. The subscript μ in W_μ^1 is not a parameter; it simply abbreviates “measure” and corresponds to the notation L_μ^1 used in [30] for the space of Radon measures which we call \mathcal{M} .

trace in W_μ^1 —here first introduced as a Radon measure on $\partial\Omega$ —is actually absolutely continuous with respect to the $(N - 1)$ -dimensional Hausdorff measure on $\partial\Omega$, and thus it can be interpreted as an element of $L^1(\partial\Omega)$.

Remark 7.2. Just like $\alpha|_{\partial\Omega}$, the outer trace β can have singular contributions, possibly even charging single points. In particular, this occurs whenever (u, α) is obtained as the weak* limit of a sequence whose gradients concentrate at a single point at the boundary, as in Example 2.13. As a side effect, the results presented here cannot be easily extended to domains with Lipschitz boundary, even with more refined tools from geometric measure theory: a normal ϱ defined \mathcal{H}^{N-1} -a.e. is simply not good enough to write (7.1) if β (and $\alpha|_{\partial\Omega}$) charges a set with Hausdorff dimension below $N - 1$.

Another simple but useful observation is that the remainder of the projection of $(u, \alpha) \in W_\mu^1$ “onto BV” given by $(u, \bar{\alpha}) \in W_\mu^1$, the *side* $(0, \chi_{\partial\Omega}\alpha) = (u, \alpha) - (u, \bar{\alpha})$ (cf. [30, Definition 4]) also is an element of W_μ^1 [30, Theorem 8]. In particular, the side of (u, α) satisfies (7.1), which reduces to

$$(7.2) \quad \int_{\partial\Omega} \varphi \varrho d(\beta - \beta^0) = \int_{\partial\Omega} \varphi d\alpha \quad \text{for every } \varphi \in C^1(\bar{\Omega}),$$

where β and β^0 denote the outer and inner traces of (u, α) , respectively. A straightforward but nonetheless remarkable consequence (also pointed out in [30, Theorem 9]) is that for every $(u, \alpha) \in W_\mu^1$, $|\alpha| = |\beta - \beta^0|$ on $\partial\Omega$ and

$$(7.3) \quad \frac{d\alpha^s}{d|\alpha^s|}(x) = \frac{d(\beta - \beta^0)^s}{d|\beta - \beta^0|^s}(x)\varrho(x) \quad \text{for } |\alpha|\text{-a.e. } x \in \partial\Omega.$$

Remark 7.3. If u is vector- instead of scalar-valued, (7.3) translates into saying that the then matrix-valued measure α^s is a matrix of rank 1 $|\alpha|$ -a.e. on $\partial\Omega$; more precisely, it is of the form $a(x) \otimes \varrho(x)$ with a suitable vector-valued $a(x)$.

Similar to BV, W_μ^1 can also be characterized using (7.1):

THEOREM 7.4 ([30, section 4]; see also [30, Definition 2]). *For any $(u, \alpha) \in W^{1,\mu}$, (7.1) is satisfied with the outer trace β of (u, α) . Conversely, for all $(u, \alpha, \beta) \in L^1(\Omega) \times \mathcal{M}(\bar{\Omega}; \mathbb{R}^N) \times \mathcal{M}(\bar{\Omega})$ such that (7.1) holds, $(u, \alpha) \in W^{1,\mu}$ and β is its outer trace.*

All of the above has obvious extensions for the vector-valued case, i.e., for the space $W_\mu^1(\bar{\Omega}; \mathbb{R}^M) \subset L^1(\Omega; \mathbb{R}^M) \times \mathcal{M}(\bar{\Omega}; \mathbb{R}^{M \times N})$ with $M > 1$. Since we defined GYM in that case, we will stick to this setting from now on. The relationship of BV, W_μ^1 and GYM can be summarized as follows:

THEOREM 7.5 (BV as a subset of W_μ^1). *We have the following chain of (natural) continuous embeddings and isomorphisms:*

$$\begin{aligned} \text{BV}(\Omega; \mathbb{R}^M) &\hookrightarrow \text{BV}(\Omega; \mathbb{R}^M) \times L^1(\partial\Omega; \mathbb{R}^M) \\ &\hookrightarrow \text{BV}(\Omega; \mathbb{R}^M) \times \mathcal{M}(\partial\Omega; \mathbb{R}^M) \cong W_\mu^1(\bar{\Omega}; \mathbb{R}^M). \end{aligned}$$

Before giving a proof, let us discuss how GYM fits into this picture. Roughly speaking, GYM exclusively contains information on gradients, unlike the other spaces appearing in Theorem 7.5: for each $\Lambda \in \text{GYM}$, the underlying deformation u is only determined up to a constant vector (Ω being connected). For a meaningful comparison, we therefore have to remove these constants, or simply drop u in the pair $(u, \alpha) \in W_\mu^1$ (recall that Du can be reconstructed from α anyway).

THEOREM 7.6 (W_μ^1 as a subset of GYM). *Let*

$$GW_\mu^1 := \{\alpha \mid (u, \alpha) \in W_\mu^1 \text{ for a suitable } u \in L^1\}.$$

(i) *For every* $\alpha \in GW_\mu^1(\bar{\Omega}; \mathbb{R}^M)$, *we have*

$$\Lambda = \Lambda(\alpha) := \left(\delta_{\nabla u(x)}, |\alpha^s|, \delta_{\frac{d\alpha^s}{d|\alpha^s|}(x)} \right) \in \text{GYM}(\bar{\Omega}; \mathbb{R}^M),$$

where ∇u denotes the density of the absolutely continuous part of α (or $Du = \alpha|_\Omega$) with respect to \mathcal{L}^N , and α^s is the singular part of α with polar decomposition $d\alpha^s(x) = \frac{d\alpha^s}{d|\alpha^s|}(x)d|\alpha^s|(x)$.

(ii) *Conversely, for every* $\Lambda \in \text{GYM}(\bar{\Omega}; \mathbb{R}^M)$, *we have an associated pair* $(u(\Lambda), \alpha(\Lambda)) \in W_\mu^1(\bar{\Omega}; \mathbb{R}^M)$, *where the underlying deformation* $u = u(\Lambda)$ *is given by Theorem 3.1 (uniquely determined up to a constant in* \mathbb{R}^M) *and* $\alpha = \alpha(\Lambda) \in GW_\mu^1(\bar{\Omega}; \mathbb{R}^M)$ *is the center of mass of* Λ , *i.e.,*

$$d\alpha(x) = \langle \nu_x, \text{id} \rangle dx + \langle \nu_x^\infty, \text{id} \rangle d\lambda(x).$$

Moreover, $\alpha(\Lambda(\gamma)) = \gamma$ for every $\gamma \in GW_\mu^1$.

Proof of Theorem 7.5. The first embedding is realized by pairing the identity and the trace operator, and the second one is trivial. Thus, we only have to show that $\text{BV}(\Omega) \times \mathcal{M}(\partial\Omega) \cong W_\mu^1(\bar{\Omega})$. Given $(u, \alpha) \in W_\mu^1(\bar{\Omega})$, we have $(u, \beta) \in \text{BV}(\Omega) \times \mathcal{M}(\partial\Omega)$, where β is the outer trace of (u, α) , and the operator $W_\mu^1(\bar{\Omega}) \rightarrow \text{BV}(\Omega) \times \mathcal{M}(\partial\Omega)$, $(u, \alpha) \mapsto (u, \beta)$, clearly is linear and continuous. It is also one-to-one, because $\alpha|_\Omega = Du$ (the derivative of $u \in \text{BV}$) and $d\alpha|_{\partial\Omega} = \varrho(x)d(\beta - \beta^0)$, where β^0 the inner trace of (u, α) which is fully determined by u and $\alpha|_\Omega = Du$.

It remains to show that $(u, \alpha) \mapsto (u, \beta)$, $W_\mu^1(\bar{\Omega}) \rightarrow \text{BV}(\Omega) \times \mathcal{M}(\partial\Omega)$ is onto. Let $(u, \beta) \in \text{BV}(\Omega) \times \mathcal{M}(\partial\Omega)$. We define $\alpha \in \mathcal{M}(\bar{\Omega}; \mathbb{R}^N)$ by $\alpha|_\Omega := Du$ and $d\alpha|_{\partial\Omega}(x) := \varrho(x)d(\beta - \beta^0)$, with the trace $\beta^0 \in L^1(\partial\Omega) \subset \mathcal{M}(\partial\Omega)$ of u in the sense of BV. Clearly, we have

$$\int_{\partial\Omega} \varphi \varrho d\beta^0 = \int_\Omega u \nabla \varphi dx + \int_\Omega \varphi dDu \text{ for every } \varphi \in C^1(\bar{\Omega}),$$

i.e., (7.1) holds for $(u, \bar{\alpha}, \beta^0) = (\beta^0)$. In addition, (7.2) holds by definition of $\alpha|_{\partial\Omega}$, and, equivalently, (7.1) holds for $(0, \chi_{\partial\Omega}\alpha, \beta - \beta^0)$. Adding up, we infer that (7.1) holds for $(u, \alpha, \beta) = (u, \bar{\alpha}, \beta^0) + (0, \chi_{\partial\Omega}\alpha, \beta - \beta^0)$. By Theorem 7.4, we conclude that $(u, \alpha) \in W_\mu^1$ and that β is its outer trace. \square

Proof of Theorem 7.6. “Moreover”: This is clear from the definition of $\Lambda(\alpha)$ and $\alpha(\Lambda)$.

(i): Given $\alpha \in GW_\mu^1$, we define $Du := \alpha|_\Omega$ (which really is the derivative of a function $u \in \text{BV}$ since $\alpha \in GW_\mu^1$), and claim that

$$\Lambda := \left(\delta_{\nabla u(x)}, |\alpha^s|, \delta_{\frac{d\alpha^s}{d|\alpha^s|}(x)} \right) \in \text{GYM}(\bar{\Omega}; \mathbb{R}^M)$$

where $\nabla u = \frac{d\alpha}{d\mathcal{L}^N} = \frac{dDu}{d\mathcal{L}^N}$. It suffices to check conditions (0)–(iii) in Theorem 3.1. By definition of Λ , its center of mass is α , and since $\alpha = Du$ in Ω , we get (0). Moreover, (i) is valid since $\langle \langle \Lambda, 1 \otimes |\cdot| \rangle \rangle = \int_\Omega |\nabla u| dx + \int_{\bar{\Omega}} d\alpha^s < \infty$, and (ii) holds with equality by definition of Λ and Du . Finally, (iii) is a consequence of the fact that $\frac{d\alpha^s}{d|\alpha^s|}(x)$ is a matrix of rank 1 of the form $a(x) \otimes \varrho(x)$ for $|\alpha^s|$ a.e. $x \in \partial\Omega$ (see Remark 7.3) combined with Lemma 2.10.

- (ii): Given $\Lambda \in \text{GYM}(\bar{\Omega}; \mathbb{R}^M)$, let $(u_n) \subset W^{1,1}(\Omega; \mathbb{R}^M)$ be a sequence such that (∇u_n) generates Λ . In particular, we have that

$$\int_{\Omega} g(x) \nabla u_n(x) \, dx \rightarrow \int_{\Omega} \langle \nu_x, \text{id} \rangle \, dx + \int_{\Omega} \langle \nu_x^{\infty}, \text{id} \rangle \, d\lambda(x)$$

for all $g \in C(\bar{\Omega})$. This means that $\nabla u_n \rightharpoonup^* \alpha$ in $\mathcal{M}(\bar{\Omega}; \mathbb{R}^{M \times N})$, where

$$d\alpha(x) := \langle \nu_x, \text{id} \rangle \, dx + \langle \nu_x^{\infty}, \text{id} \rangle \, d\lambda(x),$$

whence $\alpha \in GW_{\mu}^1$. Now let $u \in \text{BV}$ be the function given by Theorem 3.1. Since the identity $\text{id} : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}^{M \times N}$ is linear, $\pm \text{id}$ is quasiconvex and its own (generalized) recession function. Thus, inequality (ii) of Theorem 3.1 holds for $f = \text{id}$ as well as for $f = -\text{id}$, which implies equality in both cases. Hence,

$$\nabla u(x) dx = \langle \nu_x, \text{id} \rangle \, dx + \langle \nu_x^{\infty}, \text{id} \rangle \frac{d\lambda}{d\mathcal{L}^N}(x) dx,$$

and due to Theorem 3.1(0) we also have

$$dDu^s(x) = \langle \nu_x^{\infty}, \text{id} \rangle \, d\lambda^s(x) \quad \text{in } \Omega.$$

Combined, we get $dDu = \nabla u dx + dDu^s = d\alpha$ on Ω , and consequently, $(u, \alpha) \in W_{\mu}^1(\Omega; \mathbb{R}^M)$. □

8. Traces for GYM. In order to handle functionals with boundary terms, it is helpful to have a notion of trace for elements of GYM. As always, we assume that $\Omega \subset \mathbb{R}^N$ is a bounded domain with a boundary of class C^1 . Moreover, let $\Gamma \subset \partial\Omega$ be open in $\partial\Omega$. Below, Lebesgue spaces on Γ and other subsets of $\partial\Omega$ are always understood with respect to the surface measure \mathcal{H}^{N-1} .

DEFINITION 8.1 (Inner and outer trace in GYM on Γ). *Let $\Lambda \in \text{GYM}$, and let (u, α) be the associated element of the Souček space W_{μ}^1 in the sense of Theorem 7.6(ii), i.e., u is the underlying deformation of Λ (defined up to a constant), and α is the center of mass of Λ . We define the inner trace $T_i\Lambda$ of Λ in $L^1(\Gamma; \mathbb{R}^M)$ as the restriction to Γ of the inner trace of $(u, \alpha) \in W_{\mu}^1$ on $\partial\Omega$ (which coincides with the trace of u in the sense of BV). Analogously, the outer trace $T_o\Lambda \in \mathcal{M}(\Gamma; \mathbb{R}^M)$ of Λ on Γ is defined as the restriction to Γ of the outer trace of $(u, \alpha) \in W_{\mu}^1$.*

Remark 8.2.

- (i) Since the underlying deformation $u \in \text{BV}$ of Λ is only defined up to a constant in \mathbb{R}^M , the same holds for $T_i\Lambda$ and $T_o\Lambda$, with the same constant for all three. In our applications below, this does not really play a role, though, because either the constant does not matter, or one of the three is given and the constant is therefore fixed, anyway.
- (ii) If we express the derivative α of $(u, \alpha) \in W_{\mu}^1$, the element of the Souček space associated with Λ , as a generalized Young measure as in Theorem 7.6(i), we get $\bar{\Lambda}$, the center of mass of Λ . Hence $T_i\Lambda = T_i\bar{\Lambda}$ and $T_o\Lambda = T_o\bar{\Lambda}$.
- (iii) The inner trace of $(u, \alpha) \in W_{\mu}^1$ was defined as the outer trace of $(u, \bar{\alpha})$ with $\bar{\alpha} := \chi_{\Omega}\alpha$. Accordingly, we have $T_i\Lambda = T_o\Lambda^i$, where $\Lambda^i = \chi_{\Omega}\Lambda \in \text{GYM}$ is the interior part of Λ in the sense of section 4.
- (iv) The explicit formula (7.3) for the difference between outer and inner trace in the Souček space translates as follows to the trace difference in GYM:

$$(8.1) \quad dT_o\Lambda(x) - T_i\Lambda(x) d\mathcal{H}^{N-1}(x) = \langle \nu_x^{\infty}, \text{id} \rangle \varrho(x) d\lambda(x), \quad x \in \Gamma,$$

where ϱ is the outer normal on the boundary.

It is tempting to think that the traces of $\Lambda \in \text{GYM}$ should be more general objects than the traces of $(u, \alpha) \in W_\mu^1$, say, (generalized) Young measures on Γ instead of elements of $L^1(\Gamma; \mathbb{R}^M)$ or $\mathcal{M}(\Gamma; \mathbb{R}^M)$. However, as we shall see below, the inner trace enjoys a rather strong compactness property in $L^1(\Gamma; \mathbb{R}^M)$, so that a larger class is pointless, while for the outer trace, the information contained in Λ simply does not suffice to uniquely (up to a constant) determine the trace as a (generalized) Young measure on Γ .

The following theorem describes the link between traces in GYM and the limits of traces of associated generating sequences.

THEOREM 8.3. *Suppose that $(u_k) \subset \text{BV}$ is a bounded sequence generating $\Lambda = (\nu_x, \lambda, \nu_x^\infty)$, and $u_k \rightharpoonup^* u$ in BV . Then Tu_k (the trace of u_k on Γ in the sense of BV) satisfies $Tu_k \rightharpoonup^* T_o\Lambda$ in $\mathcal{M}(\Gamma; \mathbb{R}^M)$. Moreover, if $\lambda(\bar{\Gamma}) = 0$, then $Tu_k \rightarrow T_i\Lambda = T_o\Lambda = Tu$ strongly in L^1 .*

Proof. $Tu_k \rightharpoonup^* T_o\Lambda$: This immediately follows from the weak* continuity of the outer trace in the Souček space.

“Moreover”: In essence, this is closely related to the well-known fact that strict convergence in BV guarantees continuity of the trace. However, while our assumption on λ contains the essential ingredients needed for this purpose, it is weaker than strict convergence of a generating sequence, and we do not know a reference that could be directly quoted here. Therefore, we provide a short proof, although only for the case where Γ is contained in a hyperplane in \mathbb{R}^N . The general case can be recovered with standard tools in such a context, namely a decomposition of unity, maps locally straightening the boundary, and suitable error estimates. Rotating and shifting if necessary, we may even assume that $\Gamma = \{0\} \times \Gamma'$ for a suitable open $\Gamma' \subset \mathbb{R}^{N-1}$, and the outer normal on $\Gamma \subset \partial\Omega$ is given by $-e_1 = (-1, 0, \dots, 0)$.

Since $\lambda(\bar{\Gamma}) = 0$, $T_o\Lambda = T_i\Lambda$. It therefore suffices to show that (Tu_k) is Cauchy in $L^1(\Lambda; \mathbb{R}^M)$. Using the density of $C^1(\bar{\Omega}; \mathbb{R}^M)$ in $W^{1,1}(\Omega; \mathbb{R}^M)$, we may assume that $(u_k) \subset C^1(\bar{\Omega}; \mathbb{R}^M)$, while we still have that $u_k \rightharpoonup^* u$ in BV and (∇u_k) generates $\Lambda = (\nu_x, \lambda, \nu_x^\infty) \in \text{GYM}$. The basic observation now is the following: For any $w \in C^1(\bar{\Omega}; \mathbb{R}^M)$, $x' \in \Gamma'$, and $t > 0$ small enough so that $(0, t) \times \Gamma' \subset \Omega$, we have

$$(8.2) \quad \frac{1}{t} \int_0^t |w(x_1, x') - w(0, x')| dx_1 = \frac{1}{t} \int_0^t \left| \int_0^{x_1} \partial_1 w(s, x') ds \right| dx_1 \leq \int_0^t |\partial_1 w(s, x')| ds.$$

Due to (8.2),

$$\begin{aligned} & \int_{\Gamma'} |u_j(0, x') - u_k(0, x')| dx' \\ &= \int_{\Gamma'} \frac{1}{t} \int_0^t |u_j(0, x') - u_k(0, x')| dx_1 dx' \\ &= \int_{\Gamma'} \frac{1}{t} \int_0^t |u_j(0, x') - u_j(x_1, x') + u_k(x_1, x') - u_k(0, x') \\ &\quad + u_j(x_1, x') - u_k(x_1, x')| dx_1 dx' \\ &\leq \int_{\Gamma'} \frac{1}{t} \int_0^t |u_j(0, x') - u_j(x_1, x')| + |u_k(x_1, x') - u_k(0, x')| \\ &\quad + |u_j(x_1, x') - u_k(x_1, x')| dx_1 dx' \end{aligned}$$

$$\begin{aligned} &\leq \int_{\Gamma'} \frac{1}{t} \int_0^t |u_j(0, x') - u_j(x_1, x')| + |u_k(x_1, x') - u_k(0, x')| \\ &\quad + |u_j(x_1, x') - u_k(x_1, x')| \, dx_1 \, dx' \\ &\leq \int_{(0,t) \times \Gamma'} |\partial_1 u_j(x)| + |\partial_1 u_k(x)| \, dx + \frac{1}{t} \int_{(0,1) \times \Gamma'} |u_j(x) - u_k(x)| \, dx \\ &=: A(t, j, k) + \frac{1}{t} B(j, k). \end{aligned}$$

Since (u_k) is Cauchy in $L^1(\Omega; \mathbb{R}^M)$, for any $t > 0$ there is $N_0(t)$ such that $B(j, k) \leq t^2$ if $j, k \geq N_0(t)$. In addition, if $\lambda(\Gamma) = 0$,

$$\lim_{t \rightarrow 0} \lim_{k, j \rightarrow \infty} A(t, j, k) = 2 \int_{\Gamma} \langle \nu_x^\infty, |\cdot e_1| \rangle \, d\lambda(x) = 0.$$

Hence, (Tu_k) is Cauchy in $L^1(\Gamma'; \mathbb{R}^M)$. □

The weak* convergence we get for a sequence of outer traces in $\mathcal{M}(\Gamma; \mathbb{R}^M)$ is not sufficient to pass to the limit in boundary terms like an integral functional on Γ with a nonlinear integrand. Unfortunately, there seems to be no easy way to fix that, because GYM does not carry enough information to define, say, a (generalized) Young-measure outer trace:

Example 8.4. Let $\Gamma' := (0, 1)$, $(0, 1) \times \Gamma' \subset \Omega \subset \mathbb{R}^2$, and $\Gamma := \{0\} \times \Gamma' \subset \partial\Omega$ (therefore, the outer normal is $\varrho := (-1, 0)$ on Γ), and take two sequences (u_k) and (v_k) of scalar BV functions on Ω purely concentrating on Γ , with $u_k = v_0 = 0$ on $\Omega \setminus (0, 1)^2$ and the following additional properties for $(x_1, x') \in (0, 1)^2$:

- $u_k(x_1, x') = 0$ for $x_1 \geq \frac{1}{k^2}$, $\nabla u_k(x_1, x') = k^2 \varrho$ for $x_1 < \frac{1}{k^2}$. The trace is constant $u_k(0, x') = 1$ on Γ' , generating the Young measure δ_1 on Γ' as $k \rightarrow \infty$ (as well as the generalized Young measure $(\delta_1, 0, -)$).
- $v_k(x_1, x') = 0$ for $x_1 \geq \frac{1}{k^2}$, $\nabla v_k(x_1, x) = 2k^2 \varrho$ for $(x_1, x') \in (0, \frac{1}{k^2}) \times (0, \frac{1}{k})$, $\nabla v_k(x_1, x') = 0$ for $(x_1, x') \in (0, \frac{1}{k^2}) \times (\frac{1}{k}, \frac{2}{k})$, repeated $2/k$ -periodically in x' . Accordingly, the trace $v_k(0, \cdot)$ alternates between 2 and 0. In the limit as $k \rightarrow \infty$, the sequence of traces generates the Young measure $\frac{1}{2} \delta_2 + \frac{1}{2} \delta_0$ on Γ' (as well as the generalized Young measure $(\frac{1}{2} \delta_2 + \frac{1}{2} \delta_0, 0, -)$).

Despite their different Young measure “traces,” both sequences generate the *same* generalized gradient Young measure on $\bar{\Omega}$, namely $(\delta_0, \chi_\Gamma \mathcal{H}^1, \delta_\varrho)$: Concentrating at Γ with either gradient $-k^2 \varrho(x')$ on a set of measure $\frac{1}{k^2}$ or gradient $-2k^2 \varrho(x')$ on a set of measure $\frac{1}{2k^2}$ gives the same limit in GYM; the interior jumps of v_k on $(x_1, x') \in (0, \frac{1}{k}) \times (\Gamma' \cap \{\frac{j}{k} \mid j \in \mathbb{Z}\})$ are negligible, because their jump height is bounded (at most 2), and they are supported on a jump set whose surface measure is $\frac{k+1}{k^2} \rightarrow 0$.

9. Relaxation of functionals defined on $W^{1,1}$. In this section, we state and prove a relaxation result in terms of generalized Young measure suitable for functionals like the prototype in the introduction and higher-dimensional analogues. To include nonlinear boundary terms as in those examples, it would be ideal if we were able to characterize under which circumstances a pair (Λ, β) , a generalized gradient Young measure and its outer trace, can be generated by a sequence in $W^{1,1}$ such that the traces converge to β *strongly*, not just weakly* in \mathcal{M} , because this is crucial for the construction of recovery sequences even if the boundary integrand is strictly convex. But so far, we are unable to provide such a result, not even for the case $\beta \in L^1$. In particular, Example 8.4 shows that one can easily choose the “wrong” recovery sequence once oscillating traces on the boundary are possible. Of course,

none of these issues are relevant for functionals with linear boundary integrals, or no boundary integrals at all.

Below, we do allow nonlinear boundary terms, but avoid the issue with recovery sequences by imposing a few extra, possibly technical conditions on the integrand that make oscillations on the boundary energetically unfavorable, as well as concentrations at the boundary other than a simple jump growing or moving there. Some extra notation is needed for that, which we will introduce next.

9.1. Prerequisites. For $\Lambda \in \text{GYM}(\bar{\Omega}; \mathbb{R}^M)$, we define the *first moment* (sometimes also called *center of mass*) $\langle \Lambda, \text{id} \rangle \in \mathcal{M}(\bar{\Omega}; \mathbb{R}^{M \times N})$ of Λ as

$$(9.1) \quad d\langle \Lambda, \text{id} \rangle(x) := \langle \nu_x, \text{id} \rangle dx + \langle \nu_x^\infty, \text{id} \rangle d\lambda(x),$$

i.e., for any $g \in C(\bar{\Omega})$

$$\int_{\bar{\Omega}} g(x) d\langle \Lambda, \text{id} \rangle(x) = \int_{\Omega} g(x) \int_{\mathbb{R}^{M \times N}} \text{Ad}\nu_x(A) dx + \int_{\bar{\Omega}} g(x) \int_{S^{M \times N-1}} \text{Ad}\nu_x^\infty(A) d\lambda(x).$$

Alternatively, the first moment can also be written as a generalized Young measure $\bar{\Lambda} = (\bar{\nu}_x, \bar{\lambda}, \bar{\nu}_x^\infty)$, where $\bar{\nu}_x$ and $\bar{\nu}_x^\infty$ consistently are Dirac masses:

$$\bar{\nu}_x := \delta_{\langle \nu_x, \text{id} \rangle}, \quad d\bar{\lambda}(x) := |\langle \nu_x^\infty, \text{id} \rangle| d\lambda(x), \quad \bar{\nu}_x^\infty := \delta_{\frac{\langle \nu_x^\infty, \text{id} \rangle}{|\langle \nu_x^\infty, \text{id} \rangle|}}.$$

Remark 9.1. Due to Theorem 7.6(ii) and Remark 7.3, $\langle \nu_x^\infty, \text{id} \rangle$ is a rank-1 matrix of the form $a(x) \otimes \varrho(x)$ for λ -a.e. $x \in \partial\Omega$, where $\varrho(x)$ denotes the outer normal as usual and $a : \partial\Omega \rightarrow \mathbb{R}^M$ is a suitable function. As a consequence, $\Lambda \in \text{GYM}$ implies that $\bar{\Lambda} \in \text{GYM}$, by Theorem 3.1 and Lemma 2.10.

Below, we will rely on a variant of quasiconvexity at the boundary, a Jensen-type inequality we only need for recession functions:

DEFINITION 9.2 (JQCB(ϱ)). *Given a unit vector $\varrho \in S^{N-1}$ and $v^\infty : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ continuous and positively one-homogeneous, we say that $v^\infty \in \text{JQCB}(\varrho)$ if*

$$(9.2) \quad v^\infty \left(\int_{D_\varrho} \nabla \varphi dy \right) \leq \int_{D_\varrho} v^\infty(\nabla \varphi) dy \quad \text{for every } \varphi \in W_0^{1,\infty}(B; \mathbb{R}^M),$$

Remark 9.3. Clearly, all convex positively one-homogeneous functions belong to $\text{JQCB}(\varrho)$. Moreover, if v^∞ is quasiconvex at the boundary with respect to the normal ϱ as defined in [31, 26] (originally, the notion is due to [4]), i.e., for every A there exists a vector $b = b(A) \in \mathbb{R}^M$ such that

$$(9.3) \quad \int_{D_\varrho} [v^\infty(A + \nabla \varphi) - v^\infty(A)] dy \geq \int_{B \cap \{y | y \cdot \varrho = 0\}} b \cdot \varphi d\mathcal{H}^{N-1}(y) \quad \text{for every } \varphi \in W_0^{1,\infty}(B; \mathbb{R}^M),$$

then $v^\infty \in \text{JQCB}(\varrho)$. However, the proof of the latter is not entirely trivial, mainly because we cannot add nonconstant affine functions to φ in (9.3) without leaving the class of admissible test functions. Details are given in Appendix C.

LEMMA 9.4. *Let $\Lambda = (\nu_x, \lambda, \nu_x^\infty) \in \text{GYM}(\bar{\Omega}; \mathbb{R}^M)$. Then for λ -a.e. $x \in \partial\Omega$,*

$$v^\infty(\langle \nu_x^\infty, \text{id} \rangle) \leq \langle \nu_x^\infty, v^\infty \rangle \quad \text{for every } v \in \Upsilon \text{ with } v^\infty \in \text{JQCB}(\varrho(x)).$$

Proof. Fix $x_0 \in \partial\Omega$ and let $(\varphi_k) \subset W^{1,1}(B; \mathbb{R}^M)$ denote a bounded sequence such that $\text{supp } \varphi_k \subset \frac{1}{k}B$ and $(\nabla\varphi_k)$ generates $(\delta_0, \delta_0, \nu_{x_0}^\infty)$ on D_ϱ with the outer normal ϱ at x_0 (see Remark 6.5). Mollifying if necessary, we may even assume that $(\varphi_k) \subset C_c^\infty(B; \mathbb{R}^M) \subset W_0^{1,\infty}(B; \mathbb{R}^M)$. Since $v^\infty \in \text{JQCB}(\varrho)$, we have

$$v^\infty \left(\int_{D_\varrho} \nabla\varphi_k \, dy \right) \leq \int_{D_\varrho} v^\infty(\nabla\varphi_k) \, dy.$$

This implies the assertion in the limit as $k \rightarrow \infty$ because v^∞ is continuous, $\int_{D_\varrho} \nabla\varphi_k \, dy \rightarrow \langle \nu_{x_0}^\infty, \text{id} \rangle$, and $\int_{D_\varrho} v^\infty(\nabla\varphi_k) \, dy \rightarrow \langle \nu_{x_0}^\infty, v^\infty \rangle$. \square

9.2. Relaxation. Take $\Omega \in \mathbb{R}^N$ a bounded domain with a boundary of class C^1 . In addition, let $f : \bar{\Omega} \times \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ and $g : \partial\Omega \times \mathbb{R}^M \rightarrow \mathbb{R}$ be continuous, such that

- (f:1) $c^{-1}(-1 + |A|) \leq f(x, A) \leq c(1 + |A|)$ for all $x \in \bar{\Omega}$, $A \in \mathbb{R}^{M \times N}$;
- (f:2) f has a recession function in the sense of (2.14);
- (g:1) $c^{-1}(-1 + |\mu|) \leq g(x, \mu) \leq c(1 + |\mu|)$ for all $x \in \partial\Omega$, $\mu \in \mathbb{R}^M$;
- (g:2) g has a recession function in the sense of (2.14).

Here, $c > 0$ is a suitable constant. We also define

$$\begin{aligned} \Gamma_R &:= \{x \in \partial\Omega \mid g(x, \cdot) \neq 0\}, \\ \Gamma_N &:= \partial\Omega \setminus \Gamma_R. \end{aligned}$$

By definition, Γ_R and Γ_N are disjoint and cover $\partial\Omega$. Moreover, Γ_N is closed, while Γ_R is open in $\partial\Omega$. Admissible functions remain unconstrained both on Γ_R and Γ_N , which for critical points formally leads to natural boundary conditions of (nonlinear) Robin and Neumann type, respectively.

We are looking at the following problem:

$$(9.4) \quad \begin{aligned} \text{minimize } F(u) &:= \int_{\Omega} f(x, \nabla u(x)) \, dx + \int_{\partial\Omega} g(x, u) \, d\mathcal{H}^{N-1}(x) \\ &\text{subject to } u \in \mathcal{A}_C, \end{aligned}$$

where $C > 0$ is given and

$$\mathcal{A}_C := \{u \in W^{1,1}(\Omega; \mathbb{R}^M); \|u\|_{L^1(\partial\Omega; \mathbb{R}^M)} \leq C, \|\nabla u\|_{L^1(\Omega; \mathbb{R}^{M \times N})} \leq C\}.$$

Here, u as a function on $\partial\Omega$ or subsets thereof is understood in the sense of trace, and Lebesgue spaces on $\partial\Omega$ or its subsets are understood with respect to the measure \mathcal{H}^{N-1} . One particular example is the toy problem (1.1) mentioned in the introduction.

A natural extension of (9.4) to W_μ^1 is given by

$$(9.5) \quad \begin{aligned} \text{minimize } \bar{F}(u, \alpha) &:= \int_{\Omega} df(x, \alpha) + \int_{\Gamma} dg(x, \beta) \\ &\text{subject to } (u, \alpha) \in \bar{\mathcal{A}}_C, \end{aligned}$$

where β is the outer trace of (u, α) on $\partial\Omega$, β^s its singular part, and $\frac{d\beta}{d\mathcal{H}^{N-1}}$ the density of its absolutely continuous part with respect to \mathcal{H}^{N-1} ,

$$dg(x, \beta) := g \left(x, \frac{d\beta}{d\mathcal{H}^{N-1}}(x) \right) d\mathcal{H}^{N-1}(x) + g^\infty \left(x, \frac{d\beta^s}{d|\beta^s|}(x) \right) d|\beta^s|(x)$$

and

$$\bar{\mathcal{A}}_C := \{(u, \alpha) \in W^1_\mu(\Omega; \mathbb{R}^M); \|\beta\|_{\mathcal{M}(\partial\Omega; \mathbb{R}^M)} \leq C, \|\alpha\|_{\mathcal{M}(\bar{\Omega}; \mathbb{R}^{M \times N})} \leq C\}.$$

Even more generally, for $\Lambda \in \text{GYM}$ with an outer trace β on $\partial\Omega$ we may consider

$$(9.6) \quad \begin{aligned} &\text{minimize } \hat{F}(\Lambda, \beta) := \langle\langle \Lambda, f \rangle\rangle + \int_\Gamma dg(x, \beta) \\ &\text{subject to } (\Lambda, \beta) \in \hat{\mathcal{A}}_C, \end{aligned}$$

where

$$\begin{aligned} \hat{\mathcal{A}}_C &:= \{(\Lambda, \beta) \in \text{GYM}_C(\bar{\Omega}; \mathbb{R}^M) \times \mathcal{M}_C(\partial\Omega; \mathbb{R}^M) : \\ &\quad \Lambda = \bar{\Lambda} \text{ on } \Gamma, \beta \text{ is an outer trace of } \Lambda\}, \\ \text{GYM}_C &:= \{\Lambda \in \text{GYM} : \langle\langle \Lambda, 1 \otimes |\cdot| \rangle\rangle \leq C\}, \quad \mathcal{M}_C := \{\beta \in \mathcal{M} : \|\beta\|_{\mathcal{M}} \leq C\}. \end{aligned}$$

THEOREM 9.5. *Assume that in addition to (f:1), (f:2), (g:1), and (g:2), $g(z, \cdot)$ is convex, $g^\infty(z, \cdot) \geq 0$, and $f^\infty(z, \cdot) \in \text{QSLB}(\varrho(z)) \cap \text{JQCB}(\varrho(z))$ for every $z \in \Gamma_R$. Then problem (9.6) is the relaxation of both (9.4) and (9.5), i.e., $\inf F = \inf \bar{F} = \min \hat{F}$. In addition, the following holds:*

- (i) *Every minimizing sequence of either (9.4) or (9.5) contains a subsequence (not relabeled) which generates a minimizer of (9.6) in the sense that*

$$(9.7) \quad (\nabla u_k) \text{ generates } \Lambda \in \text{GYM}, \text{ and } u_k \rightharpoonup^* \beta \text{ in } \mathcal{M}(\partial\Omega; \mathbb{R}^M),$$

where on $\partial\Omega$, $u_k \in L^1(\partial\Omega; \mathbb{R}^M) \subset \mathcal{M}(\partial\Omega; \mathbb{R}^M)$ is understood in the sense of traces in $W^{1,1}$.

- (ii) *Conversely, for every $(\Lambda, \beta) = ((\nu_x, \lambda, \nu_x^\infty), \beta) \in \hat{\mathcal{A}}_C$ (arbitrary), we have $\hat{F}(\Lambda, \beta) \geq \hat{F}(\tilde{\Lambda}, \tilde{\beta})$, where $(\tilde{\Lambda}, \tilde{\beta}) = ((\tilde{\nu}_x, \tilde{\lambda}, \tilde{\nu}_x^\infty), \tilde{\beta}) \in \hat{\mathcal{A}}_C$ is given by*

$$\begin{aligned} \tilde{\Lambda} &:= \begin{cases} (\nu_x, \lambda, \nu_x^\infty) & \text{on } \Omega \cup \Gamma_N, \\ \left(- , |\langle \nu_x^\infty, \text{id} \rangle| \frac{d\lambda}{d\mathcal{H}^{N-1}} \mathcal{H}^{N-1}, \delta_{\frac{|\langle \nu_x^\infty, \text{id} \rangle|}{|\langle \nu_x^\infty, \text{id} \rangle|}} \right) & \text{on } \Gamma_R, \end{cases} \\ \tilde{\beta} &:= \begin{cases} \beta & \text{on } \Gamma_N, \\ \frac{d\beta}{d\mathcal{H}^{N-1}}(x) \mathcal{H}^{N-1} & \text{on } \Gamma_R. \end{cases} \end{aligned}$$

Moreover, if (Λ, β) is a minimizer of (9.6) (and thus also $(\tilde{\Lambda}, \tilde{\beta})$), then $(\tilde{\Lambda}, \tilde{\beta})$ is generated by a suitable minimizing sequence (u_k) of (9.4), again in the sense of (9.7).

Remark 9.6. By definition, $\tilde{\Lambda}$ satisfies $\tilde{\Lambda} = \Lambda$ on $\Omega \cup \Gamma_N$ and $\tilde{\Lambda} = \bar{\Lambda}$ on Γ_R , where $\bar{\Lambda}$ denotes the first moment of Λ as before. Analogous to Remark 9.1, it is clear that $\tilde{\Lambda}$ defined in that way really is an element of GYM for each $\Lambda \in \text{GYM}$.

Remark 9.7. Above, the sets $\mathcal{A}_C, \bar{\mathcal{A}}_C, \hat{\mathcal{A}}_C$ of admissible functions or gradient Young measures and their (outer) traces have a built-in bound, determined by the constant C . In this way, we avoid having to talk about coercivity conditions and a priori bounds. However, if a suitable a priori bound does hold, then $C > 0$ can of course be chosen large enough so that all minimizing sequences will be admissible up to a finite number of members. In this case, Theorem 9.5 also holds without the artificial bound given by C .

Proof of Theorem 9.5. Clearly, $\mathcal{A}_C \subset \overline{\mathcal{A}}_C \subset \hat{\mathcal{A}}_C$ (embedded in the sense of Theorem 7.5 and Theorem 7.6), $\bar{F} = F$ on \mathcal{A}_C , and $\hat{F} = \bar{F}$ on $\overline{\mathcal{A}}_C$. As a consequence,

$$(9.8) \quad \inf_{\mathcal{A}_C} F \geq \inf_{\overline{\mathcal{A}}_C} \bar{F} \geq \inf_{\hat{\mathcal{A}}_C} \hat{F}.$$

In addition, $\hat{\mathcal{A}}_C$ is sequentially compact with respect the convergence in (9.7), i.e., weak* convergence in $\text{GYM}(\bar{\Omega}; \mathbb{R}^M) \times \mathcal{M}(\partial\Omega; \mathbb{R}^M)$. Also observe that the first term in \hat{F} is a continuous function of Λ with respect to weak* convergence in GYM. Since g is convex and continuous, just like g^∞ , the corresponding second term in \hat{F} is a lower semicontinuous function of β with respect to weak* convergence in \mathcal{M} . In particular, we get that \hat{F} attains its minimum in $\hat{\mathcal{A}}_C$. It now suffices to show (i) and (ii), and we start with the latter.

(ii): Let $(\Lambda, \beta) \in \hat{\mathcal{A}}_C$. By definition, we also have $(\tilde{\Lambda}, \tilde{\beta}) \in \hat{\mathcal{A}}_C$. We first show that

$$(9.9) \quad \hat{F}(\Lambda, \beta) \geq \hat{F}(\tilde{\Lambda}, \tilde{\beta}).$$

Since $g \geq 0$ on Γ_R , all singular contributions on Γ_R in the boundary term are non-negative, and we get

$$\int_{\partial\Omega} dg(x, \beta) \geq \int_{\partial\Omega} dg(x, \tilde{\beta}).$$

In addition, since $f^\infty(z, \cdot) \in QSLB(\varrho(z))$ for every $z \in \Gamma_R$, we have $\langle \nu_x^\infty, f^\infty(x, \cdot) \rangle \geq 0$ for λ -a.e. $x \in \Gamma_R$, which allows us to drop any singular contribution with respect to \mathcal{H}^{N-1} on Γ_R in the first integral in (9.6):

$$\int_{\Gamma_R} \langle \nu_x^\infty, f^\infty(x, \cdot) \rangle d\lambda \geq \int_{\Gamma_R} \langle \nu_x^\infty, f^\infty(x, \cdot) \rangle \frac{d\lambda}{d\mathcal{H}^{N-1}}(x) d\mathcal{H}^{N-1}.$$

Last but not least, due to the fact that $f^\infty(z, \cdot) \in JQCB(\varrho(z))$ for every $z \in \Gamma_R$ and Lemma 9.4,

$$\begin{aligned} \int_{\Gamma_R} \langle \nu_x^\infty, f^\infty(x, \cdot) \rangle \frac{d\lambda}{d\mathcal{H}^{N-1}}(x) d\mathcal{H}^{N-1} &\geq \int_{\Gamma_R} f^\infty(x, \langle \nu_x^\infty, \text{id} \rangle) \frac{d\lambda}{d\mathcal{H}^{N-1}}(x) d\mathcal{H}^{N-1} \\ &= \int_{\Gamma_R} f^\infty \left(x, \frac{\langle \nu_x^\infty, \text{id} \rangle}{|\langle \nu_x^\infty, \text{id} \rangle|} \right) d\tilde{\lambda}, \end{aligned}$$

concluding the proof of (9.9).

In the rest of the proof, we repeatedly split generalized (gradient) Young measures into their boundary and interior parts, respectively, as defined in section 4. Furthermore, we will also use a slightly more general modification of a given generalized (gradient) Young measure; namely, for $\Lambda = (\nu_x, \lambda, \nu_x^\infty) \in \mathcal{Y}_{\text{gen}}^1$ and a Borel set $\Gamma \subset \bar{\Omega}$, with a slight abuse of notation we define

$$\chi_\Gamma \Lambda := (\chi_\Gamma(x)\nu_x + \chi_{\bar{\Omega} \setminus \Gamma} \delta_0, \lambda(\Gamma \cap \cdot), \chi_\Gamma(x)\nu_x^\infty),$$

i.e., $\chi_\Gamma \Lambda = \Lambda$ on Γ , whereas $\chi_\Gamma \Lambda$ is the “trivial” generalized Young measure $(\delta_0, 0, -)$ on $\bar{\Omega} \setminus \Gamma$. In this sense, we have in particular that $\Lambda^i = \chi_\Omega \Lambda$ and $\Lambda^b = \chi_{\partial\Omega} \Lambda$.

Now let (Λ, β) be a minimizer of (9.6). Since on Γ_R , $\tilde{\Lambda}$ coincides with the center of mass of Λ , $\chi_{\Gamma_R} \tilde{\Lambda}$ can be interpreted as an element $(0, \tilde{\alpha}) \in W_\mu^1$ according to Theorem 7.6(ii), where $d\tilde{\alpha}(x) = \chi_{\Gamma_R}(x) \langle \nu_x^\infty, \text{id} \rangle d\lambda(x)$. Since $\tilde{\lambda}|_{\Gamma_R}$ is absolutely

continuous with respect to \mathcal{H}^{N-1} by definition, the outer trace $T_o(\chi_{\Gamma_R}\tilde{\Lambda})$ is also absolutely continuous with respect to \mathcal{H}^{N-1} . Therefore, $T_o(\chi_{\Gamma_R}\tilde{\Lambda})$ is even the trace of some $w \in \text{BV}(\mathbb{R}^N \setminus \Omega; \mathbb{R}^M)$ on $\partial\Omega$, and the extension $\chi_{\mathbb{R}^N \setminus \Omega} w$ of w to \mathbb{R}^N is an element of $\text{BV}(\mathbb{R}^N; \mathbb{R}^M)$. Consequently, we can choose a bounded sequence $(w_k) \subset W^{1,1}(\mathbb{R}^N; \mathbb{R}^M)$ such that $w_k = w$ on $\mathbb{R}^N \setminus \Omega$, $w_k \rightarrow \chi_{\mathbb{R}^N \setminus \Omega} w$ strictly in $\text{BV}(\mathbb{R}^N; \mathbb{R}^M)$, $w_k = 0$ on $\Omega \setminus (\partial\Omega)_{\frac{1}{k}}$. In particular, $(\nabla w_k) \subset L^1(\Omega; \mathbb{R}^{M \times N})$ generates $\chi_{\Gamma_R}\tilde{\Lambda}^b = \chi_{\Gamma_R}\tilde{\Lambda}$. This also implies that $\|w_k\|_{W^{1,1}} \rightarrow \|\chi_{\Gamma_R}\tilde{\Lambda}^b\|_{\mathcal{Y}_{\text{gen}}^1}$, and, modifying w_k by multiplying with a suitable sequence of constants converging to 1, we may assume w.l.o.g. that $\|w_k\|_{W^{1,1}} \leq \|\chi_{\Gamma_R}\tilde{\Lambda}^b\|_{\mathcal{Y}_{\text{gen}}^1}$ for all k .

Now take two other bounded sequences $(y_k), (z_k) \in W^{1,1}$ such that (∇y_k) generates $\chi_{\Gamma_N}\tilde{\Lambda}^b = \chi_{\Gamma_N}\tilde{\Lambda}$ and (∇z_k) generates $\Lambda^i = \Lambda^i = \chi_{\Omega}\tilde{\Lambda}$. In addition, w.l.o.g., we may assume that $\|y_k\|_{W^{1,1}} \leq \langle \langle \chi_{\Gamma_N}\tilde{\Lambda}^b, 1 \otimes |\cdot| \rangle \rangle$ and $\|z_k\|_{W^{1,1}} \leq \langle \langle \tilde{\Lambda}^i, 1 \otimes |\cdot| \rangle \rangle$. For $u_k := z_k + y_k + w_k$, we get that (∇u_k) generates $\tilde{\Lambda}$ due to Lemma 2.16. As a consequence, $u_k \rightharpoonup^* \tilde{\beta} = T_o\tilde{\Lambda}$ on $\partial\Omega$.

To see that (u_k) is a minimizing sequence for (9.4), observe the following: $z_k \rightarrow T_i\tilde{\Lambda}$ strongly in L^1 on $\partial\Omega$ by Theorem 8.3, $y_k \rightarrow 0$ strongly in L^1 on every compact $K \subset \Gamma_R$, and $w_k = \tilde{\gamma} := T_o(\chi_{\Gamma_R}\tilde{\Lambda}) = \chi_{\Gamma_R}(T_o\tilde{\Lambda} - T_i\tilde{\Lambda})$ is a constant sequence on Γ_R . In particular, besides weak* convergence in \mathcal{M} , we also have

$$u_k \rightarrow T_i\tilde{\Lambda} + \tilde{\gamma} = \tilde{\beta} \text{ strongly in } L^1_{\text{loc}}(\Gamma_R; \mathbb{R}^M),$$

and since both g and g^∞ are continuous with $g(x, \cdot) = 0$ for $x \in \Gamma_N = \partial\Omega \setminus \Gamma_R$, we conclude that

$$\int_{\Gamma_R} g(x', u_k(x')) \, d\mathcal{H}^{N-1}(x') \rightarrow \int_{\Gamma_R} g(x', \tilde{\beta}(x')) \, d\mathcal{H}^{N-1}(x').$$

Altogether, we have $F(u_k) \rightarrow \hat{F}(\tilde{\Lambda}, \tilde{\beta})$ as $k \rightarrow \infty$.

(i): Let (u_k) be a minimizing sequence of (9.4), which generates $(\Lambda_0, \beta_0) \in \hat{\mathcal{A}}_C$. Due to lower semicontinuity, $\hat{F}(\Lambda_0, \beta_0) \leq \inf_{\mathcal{A}_C} F = \lim F(u_k)$. Picking a minimizer $(\tilde{\Lambda}, \tilde{\beta})$ as in (ii), we get another minimizing sequence (w_k) such that $\inf_{\mathcal{A}_C} F = \lim F(w_k) = \hat{F}(\tilde{\Lambda}, \tilde{\beta}) = \min_{\hat{\mathcal{A}}_C} \hat{F}$. As a consequence, $\min_{\hat{\mathcal{A}}_C} \hat{F} \leq \hat{F}(\Lambda_0, \beta_0) \leq \inf_{\mathcal{A}_C} F = \min_{\hat{\mathcal{A}}_C} \hat{F}$, and we have equality everywhere. \square

Appendix A. Proof of Proposition 2.8.

We start with the following lemma.

LEMMA A.1 (Special case of [5, Proposition 5.5]). *Assume that $f = g \otimes v$ with $(g, v) \in C(\bar{\Omega}) \times \Upsilon$ and that F is given by (2.10). Let $\mathcal{U} \subset \text{BV}(\Omega; \mathbb{R}^N)$ be an additively closed set, and suppose that $F : \mathcal{U} \rightarrow \mathbb{R}$ is bounded from below. Then F is weak* lower semicontinuous along all sequences $(c_n)_{n \in \mathbb{N}} \subset \mathcal{U}$ that are bounded in $\text{BV}(\Omega; \mathbb{R}^N)$ and in addition satisfy that $S_n := \{c_n \neq 0\} \subset (\partial\Omega)_{r_n}$ for a decreasing sequence $r_n \searrow 0$.*

Remark A.2. The class of integrands f for which [5, Proposition 5.5] is stated and proved is larger than the product form we assume in Lemma A.1 above. Notice that this product form automatically has a recession function in the very strong sense of (2.14), which corresponds to the crucial assumption (f:2) in [5]. For our purposes here, integrands in product form are enough.

Proof of Proposition 2.8. For fixed $\varepsilon > 0$, we cover $\partial\Omega$ by the following collection of balls:

$$(A.1) \quad \partial\Omega \subset \bigcup_{x \in \partial\Omega} \bigcup_{\delta \leq \tilde{\delta}(x, \varepsilon)} B_\delta(x),$$

where $\tilde{\delta}(x, \varepsilon)$ is any such radius such that the condition characterizing quasisublinear growth from below as stated in Definition 2.5 holds; here we recall that if this condition holds with the ball of radius $\tilde{\delta}(x, \varepsilon)$ it also holds for any ball of smaller radius.

Furthermore, since $\partial\Omega$ is compact we can select from the cover in (A.1) a finite subcover

$$\partial\Omega \subset \bigcup_{j=1}^J B_{\delta_j}(x_j)$$

with the radii bounded from below, i.e., $\delta_j \geq \delta_0$ for some $\delta_0 = \delta_0(\varepsilon)$. In fact, since the balls $B_{\delta_j}(x_j)$ are open and the collection is finite, we can even find $\alpha > 0$ such that balls of the radii $\delta_j - \alpha$ still cover $\partial\Omega$, i.e.,

$$\partial\Omega \subset \bigcup_{j=1}^J \overline{B_{\delta_j - \alpha}(x_j)}.$$

Let us now apply the local decomposition Lemma 2.2 to the sequence $(c_n)_{n \in \mathbb{N}}$ with the compact sets

$$\begin{aligned} K_1 &= \overline{B_{\delta_1 - \alpha}(x_1)} \cap \overline{\Omega}, \\ &\vdots \\ K_J &= \overline{B_{\delta_J - \alpha}(x_J)} \cap \overline{\Omega}, \\ K_{J+1} &= \Omega \setminus \overline{\bigcup_{j=1}^J B_{\delta_j - \alpha}(x_j)}; \end{aligned}$$

so we can write

$$(A.2) \quad c_n = c_{1,n} + c_{2,n} + \dots + c_{J+1,n},$$

where $c_{j,n}$ are supported in $B_{\delta_j}(x_j)$ for $j = 1, \dots, J$ is and $c_{J+1,n}$ is supported in Ω . Notice that we need n large enough depending on α and δ_0 in order to fulfill these requirements. Moreover, $c_{1,n}, \dots, c_{J,n}$ retain the property of the original sequence to be concentrating on the boundary while $c_{J+1,n} = 0$ for large n , and so

$$(A.3) \quad F(0) = \lim_{n \rightarrow \infty} F(c_{J+1,n}).$$

As the next step, we define the auxiliary functionals

$$G_j(u) := \int_{\Omega \cap B_{\delta_j}(x_j)} f(x, \nabla u(x)) + \varepsilon |\nabla u(x)| \, dx,$$

$$u \in \mathcal{U}_j := \{w \in \text{BV}(\Omega \cap B_{\delta_j}(x_j); \mathbb{R}^n) \text{ with } w = 0 \text{ in a neighborhood of } \partial B_{\delta_j}(x_j)\}.$$

Each is bounded from below due to the given quasisublinear growth from below; cf. Definition 2.5. Therefore, they are lower semicontinuous along sequences purely concentrating on the boundary due to Lemma A.1; in particular, G_j is lower semicontinuous along $(c_{j,n})$ (note that indeed $(c_{j,n})$ vanishes near $\partial B_{\delta_j}(x_j)$). As a consequence,

$$\begin{aligned} \lim_{n \rightarrow \infty} F(c_{j,n}) - F(0) &= \lim_{n \rightarrow \infty} G_j(c_{j,n}) - G(0) - \varepsilon \int_{\Omega \cap B_{\delta_j}(x_j)} |\nabla c_n(x)| \, dx \\ &\geq -\varepsilon \lim_{n \rightarrow \infty} \int_{\Omega \cap B_{\delta_j}(x_j)} |\nabla c_n(x)| \, dx. \end{aligned}$$

By (A.3) and Proposition 2.3 (which applies to F as well as to $u \mapsto |\nabla u|$), the sum over j yields that

$$(A.4) \quad \lim_{n \rightarrow \infty} F(c_n) - F(0) \geq -\varepsilon \lim_{n \rightarrow \infty} \int_{\Omega} |\nabla c_n(x)| \, dx,$$

As $\varepsilon > 0$ is arbitrary and $(c_n)_n$ is bounded in $W^{1,1}(\Omega; \mathbb{R}^M)$, the claim follows.

The following appendix links together generalized Young measures and the so-called DiPerna–Majda measures [10] which allow for much more general treatment of concentration effects and which were used by the last two authors in many occasions [8, 12, 15, 22, 23]. We confine ourselves to a special case of $p = 1$ and of the compactification of $\mathbb{R}^{M \times N}$ by the sphere when one can show a one-to-one correspondence of the DiPerna–Majda and generalized Young measures. More general integrands are treated with this tool in [22, 24, 29].

Appendix B. Relation of generalized Young measures to DiPerna–Majda measures. A different but equivalent description of oscillation and concentration effects in L^p -bounded sequences can be reached by means of DiPerna–Majda measures [10, 24, 25]. However, DiPerna–Majda measures allow for much more general test functions if we replace the sphere used in the definition of $\mathcal{Y}_{\text{gen}}^1$ by a larger set. We refer to [29] for a comprehensive treatment.

In what follows we will work mostly with a particular compactification of $\mathbb{R}^{M \times N}$, namely, with the compactification by the sphere. We will consider the following ring of continuous bounded functions:

(B.1)

$$\mathcal{S} := \left\{ v_0 \in C(\mathbb{R}^{M \times N}) : \text{there exist } c \in \mathbb{R}, v_{0,0} \in C_0(\mathbb{R}^{M \times N}), \right. \\ \left. \text{and } v_{0,1} \in C(S^{M \times N - 1}) \text{ s.t.} \right. \\ \left. v_0(A) = c + v_{0,0}(A) + v_{0,1} \left(\frac{A}{|A|} \right) \frac{|A|}{1 + |A|} \text{ if } A \neq 0 \text{ and } v_0(0) = v_{0,0}(0) + c \right\},$$

where $S^{M \times N - 1}$ denotes the $(MN - 1)$ -dimensional unit sphere in $\mathbb{R}^{M \times N}$. Then $\beta_S \mathbb{R}^{M \times N}$ is homeomorphic to the unit ball $\overline{B}(0, 1) \subset \mathbb{R}^{M \times N}$ via the mapping $d : \mathbb{R}^{M \times N} \rightarrow \overline{B}(0, 1)$, $d(A) := A/(1 + |A|)$ for all $A \in \mathbb{R}^{M \times N}$. Note that $d(\mathbb{R}^{M \times N})$ is dense in $\overline{B}(0, 1)$. Notice also that, in view of (2.4),

$$\Upsilon = \{v : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}; v/(1 + |\cdot|) \in \mathcal{S}\}.$$

Let $\sigma \in \mathcal{M}(\overline{\Omega})$ be a positive Radon measure on a bounded domain $\Omega \subset \mathbb{R}^N$. A mapping $\hat{\nu} : x \mapsto \hat{\nu}_x$ belongs to the space $L_w^\infty(\overline{\Omega}, \sigma; \mathcal{M}(\beta_S \mathbb{R}^{M \times N}))$ if it is weakly* σ -measurable (i.e., for any $v_0 \in C_0(\mathbb{R}^{M \times N})$, the mapping $\overline{\Omega} \rightarrow \mathbb{R} : x \mapsto \int_{\beta_S \mathbb{R}^{M \times N}} v_0(A) d\hat{\nu}_x(A)$ is σ -measurable in the usual sense). If additionally $\hat{\nu}_x \in \mathcal{M}_1^+(\beta_S \mathbb{R}^{M \times N})$ for σ -a.a. $x \in \overline{\Omega}$, the collection $\{\hat{\nu}_x\}_{x \in \overline{\Omega}}$ is the so-called Young measure on $(\overline{\Omega}, \sigma)$ [35]; see also [3, 29, 33, 34].

DiPerna and Majda [10] showed that having a bounded sequence in $(Y_k)_{k \in \mathbb{N}} \subset L^1(\Omega; \mathbb{R}^{M \times N})$ with $1 \leq p < +\infty$ and Ω an open domain in \mathbb{R}^N , there exists a subsequence (denoted by the same indices), a positive Radon measure $\sigma \in \mathcal{M}(\overline{\Omega})$, and a Young measure $\hat{\nu} : x \mapsto \hat{\nu}_x$ on $(\overline{\Omega}, \sigma)$ such that $(\sigma, \hat{\nu})$ is attainable by this subsequence in the sense that for all $g \in C(\overline{\Omega})$ and all $v_0 \in \mathcal{S}$,

$$(B.2) \quad \lim_{k \rightarrow \infty} \int_{\Omega} g(x) v(Y_k(x)) dx = \int_{\Omega} \int_{\beta_S \mathbb{R}^{M \times N}} g(x) v_0(A) d\hat{\nu}_x(A) d\sigma(x),$$

where $v \in \Upsilon$. In particular, putting $v_0 := v/(1 + |\cdot|) = 1$ in (B.2) we can see that

$$(B.3) \quad \lim_{k \rightarrow \infty} (1 + |Y_k|) = \sigma \quad \text{weakly* in } \mathcal{M}(\bar{\Omega}).$$

If (B.2) holds, we say that $\{y_k\}_{k \in \mathbb{N}}$ generates $(\sigma, \hat{\nu})$ and we denote the set of all such pairs $(\sigma, \hat{\nu})$ by DM.

Comparing (B.3) and (2.13) applied for $v(A) := |A|$ we immediately see that

$$(B.4) \quad \sigma = \left(\int_{\mathbb{R}^{M \times N}} (1 + |A|) d\nu_x(A) \right) \mathcal{L}^N + \lambda.$$

In particular, $\sigma_s = \lambda_s$, i.e., singular parts (in the Lebesgue decomposition) of both measures coincide. In view of [12, (A.7)], for all $g \in C(\bar{\Omega})$ and all $v \in \Upsilon_{\mathcal{S}}(\mathbb{R}^{M \times N})$

$$(B.5) \quad \begin{aligned} \lim_{k \rightarrow \infty} \int_{\Omega} g(x) v(y_k(x)) dx &= \int_{\Omega} \int_{\mathbb{R}^{M \times N}} g(x) v(A) d\nu_x(A) dx \\ &+ \int_{\bar{\Omega}} \int_{\beta_{\mathcal{S}} \mathbb{R}^{M \times N} \setminus \mathbb{R}^{M \times N}} g(x) \frac{v^{\infty}(A)}{1 + |A|} d\hat{\nu}_x(A) d\sigma(x). \end{aligned}$$

Comparing this with (2.13), we get

$$(B.6) \quad \int_{\bar{\Omega}} \int_{\beta_{\mathcal{S}} \mathbb{R}^{M \times N} \setminus \mathbb{R}^{M \times N}} g(x) \frac{v^{\infty}(A)}{1 + |A|} d\hat{\nu}_x(A) d\sigma(x) = \int_{\bar{\Omega}} \int_{S^{M \times N-1}} v^{\infty}(A) \nu_x^{\infty}(dA) g(x) d\lambda(x).$$

Setting $v^{\infty}(A) := |A|$ in the above identity (B.6) we get that

$$(B.7) \quad \frac{d\lambda}{d\sigma}(x) = \hat{\nu}_x(\beta_{\mathcal{S}} \mathbb{R}^{M \times N} \setminus \mathbb{R}^{M \times N}).$$

Substituting back to (B.6) we finally obtain that for σ -a.a. $x \in \bar{\Omega}$ and all continuous and positively one-homogeneous functions v^{∞}

$$(B.8) \quad \int_{\beta_{\mathcal{S}} \mathbb{R}^{M \times N} \setminus \mathbb{R}^{M \times N}} \frac{v^{\infty}(A)}{1 + |A|} d\hat{\nu}_x(A) = \hat{\nu}_x(\beta_{\mathcal{S}} \mathbb{R}^{M \times N} \setminus \mathbb{R}^{M \times N}) \int_{S^{M \times N-1}} v^{\infty}(A) \nu_x^{\infty}(dA).$$

Comparing (B.2) and (B.5) we get for all $v \in \Upsilon$ and almost all $x \in \Omega$

$$(B.9) \quad \frac{d\mathcal{L}^N}{d\sigma}(x) \int_{\mathbb{R}^{M \times N}} v(A) d\nu_x(A) = \int_{\mathbb{R}^{M \times N}} \frac{v(A)}{1 + |A|} d\hat{\nu}_x(A),$$

which connects the Young measure ν and the restriction of $\hat{\nu}$ on $\mathbb{R}^{M \times N}$. Altogether, if $(\nu, \lambda, \nu^{\infty}) \in \mathcal{Y}_{\text{gen}}^1$ we construct $(\sigma, \hat{\nu}) \in \text{DM}$ from (B.4), (B.7), (B.8), and from (B.9) and similarly, given $(\sigma, \hat{\nu}) \in \text{DM}$, we construct $(\nu, \lambda, \nu^{\infty}) \in \mathcal{Y}_{\text{gen}}^1$.

Appendix C. Quasiconvexity at the boundary versus JQCB. We recall the following notion originally introduced in [4], in the form given in [31, 26]:

DEFINITION C.1 (qcb: quasiconvex at the boundary). *Given $v \in C(\mathbb{R}^{M \times N})$, we say that v is quasiconvex at the boundary (qcb) at the matrix $A \in \mathbb{R}^{M \times N}$ with respect to the normal $\varrho \in S^{N-1}$ if there exists a vector $b = b(A) \in \mathbb{R}^M$ such that*

$$(C.1) \quad \int_{D_{\varrho}} [v(A + \nabla \varphi) - v(A)] dy \geq \int_{B \cap \{y \cdot \varrho = 0\}} b \cdot \varphi d\mathcal{H}^{N-1}(y) \text{ for every } \varphi \in W_0^{1,\infty}(B; \mathbb{R}^M).$$

Remark C.2.

- (i) The half-ball $D_\varrho = B \cap \{y \mid y \cdot \varrho < 0\}$ is a prototypical example for a so-called “standard boundary domain” with respect to ϱ . Standard boundary domains are characterized by the fact that their boundary contains a planar part perpendicular to ϱ , and there the test functions φ remain unconstrained, while they have to vanish on the rest of the boundary. Quasiconvexity at the boundary can be equivalently defined using another such domain instead of D_ϱ [31, 26].
- (ii) The boundary integral on the right-hand side of (C.1) can be rewritten as a volume integral, using the Gauss theorem and the boundary conditions for φ :

$$(C.2) \quad \int_{B \cap \{y \mid y \cdot \varrho = 0\}} b \cdot \varphi \, d\mathcal{H}^{N-1}(y) = \int_{D_\varrho} (b \otimes \varrho) : D\varphi \, dy.$$

- (iii) Suppose that v has a recession function v^∞ . Then, if v is qcb at every $A \in \mathbb{R}^{M \times N}$ with respect to ϱ , so is v^∞ .

The link to JQCB(ϱ) as introduced in Definition 9.2 is the following:

LEMMA C.3. *Let $\varrho \in S^{N-1}$, and let $v^\infty : \mathbb{R}^{M \times N} \rightarrow \mathbb{R}$ be continuous and positively one-homogeneous. If v^∞ is qcb at A with respect to ϱ for every $A \in \mathbb{R}^{M \times N}$, then $v^\infty \in \text{JQCB}(\varrho)$, i.e.,*

$$(C.3) \quad v^\infty \left(\int_{D_\varrho} \nabla \varphi \, dy \right) \leq \int_{D_\varrho} v^\infty(\nabla \varphi) \, dy \quad \text{for every } \varphi \in W_0^{1,\infty}(B; \mathbb{R}^M).$$

Proof. Let $Q = Q(\varrho) \subset \mathbb{R}^N$ be an open unit cube centered at the origin, such that one of the sides of Q is perpendicular to ϱ , and let

$$Q_\varrho := Q \cap H_\varrho, \quad \text{where } H_\varrho := \{y \in \mathbb{R}^N \mid y \cdot \varrho < 0\}.$$

Q_ϱ is another example for a standard boundary domain with respect to ϱ , and it is not difficult to check (see [31, 26] for details) that for $v = v^\infty$, (C.1) is equivalent to

$$(C.4) \quad \int_{Q_\varrho} [v^\infty(A + \nabla \varphi) - v^\infty(A)] \, dy \geq \int_{Q_\varrho} (b \otimes \varrho) : \nabla \varphi \, dy \quad \text{for every } \varphi \in W_0^{1,\infty}(Q; \mathbb{R}^M),$$

where we also used the analogue of (C.2) on Q_ϱ . Moreover, the class of test functions φ in (C.4) can be extended to BV by a density argument. In particular, using the density of $W^{1,\infty}$ in BV with respect to strict convergence, we get

$$(C.5) \quad \int_{H_\varrho} [dv^\infty(A + D\varphi)(y) - v^\infty(A) \, dy] \geq \int_{H_\varrho} (b \otimes \varrho) : dD\varphi(y)$$

for every $\varphi \in \text{BV}(\mathbb{R}^N; \mathbb{R}^M)$ with $\varphi = 0$ on $\mathbb{R}^N \setminus \bar{Q}$.

Finally, since $dv^\infty(\tilde{A} + D\varphi)(y) - v^\infty(\tilde{A}) \, dy = 0$ on $\mathbb{R}^N \setminus \text{supp } \varphi$ for all $\tilde{A} \in \mathbb{R}^{M \times N}$, we may replace the fixed matrix A by any measurable function $F : \mathbb{R}^N \rightarrow \mathbb{R}^{M \times N}$ such that $F(y) = A$ for all $y \in H_\varrho \cap \text{supp } \varphi$:

$$(C.6) \quad \int_{H_\varrho} [dv^\infty(F + D\varphi)(y) - v^\infty(F(y)) \, dy] \geq \int_{H_\varrho} (b \otimes \varrho) : dD\varphi(y)$$

for every $\varphi \in \text{BV}(\mathbb{R}^N; \mathbb{R}^M)$ with $\varphi = 0$ on $\mathbb{R}^N \setminus \bar{Q}$.

Analogous to the equivalence of (C.1) and (C.4), our assertion (C.3) holds if and only if for every $\psi \in W_0^{1,\infty}(Q; \mathbb{R}^M)$,

$$(C.7) \quad v^\infty \left(\int_{H_\varrho} \nabla \psi \, dy \right) \leq \int_{H_\varrho} v^\infty(\nabla \psi) \, dy.$$

Therefore, given $\psi \in W_0^{1,\infty}(Q; \mathbb{R}^M)$, let us see that (C.7) indeed holds.

Due to the boundary conditions of ψ , $\int_{Q_\varrho} (\nabla \Psi(y)) \varrho^\perp \, dy = 0$ for every vector $\varrho^\perp \in \mathbb{R}^N$ perpendicular to ϱ . Therefore, there exists a vector $a = a(\psi) \in \mathbb{R}^M$ such that

$$a \otimes \varrho = \int_{Q_\varrho} \nabla \Psi(y) \, dy \in \mathbb{R}^{M \times N}.$$

We will use (C.6) with the following functions: For $y \in Q_\varrho$ and $k \geq 2$, define $F_k \in L^\infty(Q_\varrho; \mathbb{R}^{M \times N})$ and $\varphi_k \in \text{BV}(H_\varrho; \mathbb{R}^M)$ such that

$$F_k(y) := \begin{cases} 0 & \text{for } -1 < \varrho \cdot y < -\frac{1}{k}, \\ k(a \otimes \varrho) & \text{for } -\frac{1}{k} \leq \varrho \cdot y < 0, \end{cases} \quad \begin{aligned} \nabla \varphi_k(y) &:= -F_k(y) + \nabla \psi_k(y), \\ &\text{with } \psi_k(y) := k^{N-1} \psi(ky), \end{aligned}$$

extended by zero to all of \mathbb{R}^N . Here, notice that $F_k(y) = \nabla f_k(y)$, where ∇f_k is the absolutely continuous part of the BV derivative $Df_k(y)$ of a potential $f_k \in \text{BV}(\mathbb{R}^N; \mathbb{R}^M)$ such that

$$\begin{aligned} f_k &= 0 \text{ on } \mathbb{R}^N \setminus \overline{Q_\varrho}, \\ f_k &\text{ is piecewise affine and continuous in } Q_\varrho \text{ with} \\ f_k &= 0 \text{ on } \{y \in Q_\varrho \mid \varrho \cdot y < -\frac{1}{k}\} \text{ and} \\ \nabla f_k &= k(a \otimes \varrho) \text{ on } \{y \in Q_\varrho \mid -\frac{1}{k} < \varrho \cdot y < 0\}. \end{aligned}$$

In particular, $\varphi_k := -f_k + \psi_k \in \text{BV}(\mathbb{R}^N; \mathbb{R}^M)$ is admissible as a test function in (C.6), and F_k is constant on $\text{supp } \varphi_k \cap H_\varrho \subset (\frac{1}{k}Q) \cap H_\varrho$ as required for (C.6). Consequently, the singular part of Df_k is supported on the jump set of f_k given by

$$J_k := \{y \in \partial Q_\varrho \mid -\frac{1}{k} \leq \varrho \cdot y \leq 0\} \cup (\partial Q_\varrho \cap \partial H_\varrho),$$

$f_k(\mathbb{R}^N) \subset [0, a]$ (the closed one-dimensional line segment connecting 0 and a in \mathbb{R}^M), and

$$(C.8) \quad |Df_k|(J_k \cap H_\varrho) \leq |a| \mathcal{H}^{N-1}(J_k \cap H_\varrho) = |a| \frac{2N}{k} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Moreover,

$$\int_{H_\varrho} dD\varphi_k(y) = - \int_{J_k \cap H_\varrho} dDf_k(y) - \int_{H_\varrho} F_k \, dy + \int_{H_\varrho} \nabla \psi_k \, dy = 0 - a \otimes \varrho + a \otimes \varrho = 0.$$

Therefore, the right-hand side of (C.6) with $F = F_k$ and $\varphi = \varphi_k$ vanishes, and we obtain

$$(C.9) \quad \int_{H_\varrho} [dv^\infty(F_k + D\varphi_k)(y) - v^\infty(F_k(y))] \, dy \geq 0.$$

Using first the definition of φ_k and then the definition of F_k , a change of variables $z = ky$, and the homogeneity of v^∞ , we infer that

$$\begin{aligned} (C.10) \quad 0 &\leq \int_{J_k \cap H_\varrho} dv^\infty(Df_k)(y) + \int_{H_\varrho} [v^\infty(\nabla\psi_k(y)) - v^\infty(F_k(y))] dy \\ &= \int_{J_k \cap H_\varrho} dv^\infty(Df_k)(y) + \int_{H_\varrho} v^\infty(\nabla\psi(z)) dz - v^\infty(a \otimes \varrho). \end{aligned}$$

Since $\int_{Q_\varrho} \nabla\psi dy = a \otimes \varrho$, due to (C.8), passing to the limit as $k \rightarrow \infty$ we finally get (C.7). \square

Acknowledgment. This research was partly conducted when M.K. held the visiting Giovanni-Prodi professorship in the Institute of Mathematics, University of Würzburg. Its support and hospitality are gratefully acknowledged.

REFERENCES

- [1] J. J. ALIBERT AND G. BOUCHITTÉ, *Non-uniform integrability and generalized Young measures*, J. Convex Anal., 4 (1997), pp. 125–145.
- [2] L. AMBROSIO, N. FUSCO, AND D. PALLARA, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Math. Monogr., Clarendon Press, Oxford, 2000.
- [3] J. M. BALL, *A version of the fundamental theorem for Young measures*, in PDEs and Continuum Models of Phase Transition, M. Rascle, D. Serre, M. Slemrod, eds., Lect. Notes Phys. 344, Springer, Berlin, 1989, pp. 207–215.
- [4] J. M. BALL AND J. MARSDEN, *Quasiconvexity at the boundary, positivity of the second variation and elastic stability*, Arch. Ration. Mech. Anal., 86 (1984), pp. 251–277.
- [5] B. BENEŠOVÁ, S. KRÖMER, AND M. KRUŽÍK, *Boundary effects and weak* lower semicontinuity for signed integral functionals on BV*, ESAIM: Control Optim. Calc. Var., 21 (2015), pp. 513–534.
- [6] B. BENEŠOVÁ AND M. KRUŽÍK, *Weak lower semicontinuity of integral functionals and applications*, SIAM Rev., 59 (2017), pp. 703–766.
- [7] M. BILDHAUER, *Convex Variational Problems: Linear, Nearly Linear and Anisotropic Growth Conditions*, Lect. Notes Math. 1818, Springer, Berlin, 2008.
- [8] M. CLAEYS, D. HENRION, AND M. KRUŽÍK, *Semi-definite relaxations for optimal control problems with oscillation and concentration effects*, ESAIM: Control Optim. Calc. Var., 23 (2017), pp. 95–117.
- [9] B. DACOROGNA, *Direct Methods in the Calculus of Variations*, 2nd ed., Springer, Berlin, 2008.
- [10] R. J. DIPIERNA AND A. J. MAJDA, *Oscillations and concentrations in weak solutions of the incompressible fluid equations*, Commun. Math. Phys., 108 (1987), pp. 667–689.
- [11] I. FONSECA, *Lower semicontinuity of surface energies*, Proc. Roy. Soc. Edinburgh, 120A (1992), pp. 95–115.
- [12] I. FONSECA AND M. KRUŽÍK, *Oscillations and concentrations generated by A-free mappings and weak lower semicontinuity of integral functionals*, ESAIM Control Optim. Calc. Var., 16 (2010), pp. 472–502.
- [13] I. FONSECA, S. MÜLLER, AND P. PEDREGAL, *Analysis of concentration and oscillation effects generated by gradients*, SIAM J. Math. Anal., 29 (1998), pp. 736–756.
- [14] A. KAŁAMAJSKA, S. KRÖMER, AND M. KRUŽÍK, *Sequential weak continuity of null Lagrangians at the boundary*, Calc. Var., 49 (2014), pp. 1263–1278.
- [15] A. KAŁAMAJSKA AND M. KRUŽÍK, *Oscillations and concentrations in sequences of gradients*, ESAIM Control Optim. Calc. Var., 14 (2008), pp. 71–104.
- [16] D. KINDERLEHRER AND P. PEDREGAL, *Characterization of Young measures generated by gradients*, Arch. Ration. Mech. Anal., 115 (1991), pp. 329–365.
- [17] D. KINDERLEHRER AND P. PEDREGAL, *Gradient Young measures generated by sequences in Sobolev spaces*, J. Geom. Anal., 4 (1994), pp. 59–90.
- [18] B. KIRCHHEIM AND J. KRISTENSEN, *Automatic convexity of rank-1 convex functions*, C. R., Math. Acad. Sci. Paris, 349 (2011), pp. 407–409.
- [19] B. KIRCHHEIM AND J. KRISTENSEN, *On rank one convex functions that are homogeneous of degree one*, Arch. Ration. Mech. Anal., 221 (2016), pp. 527–558.

- [20] J. KRISTENSEN AND F. RINDLER, *Characterization of generalized gradient Young measures generated by sequences in $W^{1,1}$ and BV* , Arch. Ration. Mech. Anal., 197 (2010), pp. 539–598; and Erratum, 203 (2012), pp. 693–700.
- [21] S. KRÖMER, *On the role of lower bounds in characterizations of weak lower semicontinuity of multiple integrals*, Adv. Calc. Var., 3 (2010), pp. 378–408.
- [22] S. KRÖMER AND M. KRUŽÍK, *Oscillations and concentrations in sequences of gradients up to the boundary*, J. Convex Anal., 20 (2013), pp. 723–752.
- [23] M. KRUŽÍK, *Quasiconvexity at the boundary and concentration effects generated by gradients*, ESAIM Control Optim. Calc. Var., 19 (2013), pp. 679–700.
- [24] M. KRUŽÍK AND T. ROUBÍČEK, *On the measures of DiPerna and Majda*, Math. Bohem., 122 (1997), pp. 383–399.
- [25] M. KRUŽÍK AND T. ROUBÍČEK, *Optimization problems with concentration and oscillation effects: relaxation theory and numerical approximation*, Numer. Funct. Anal. Optim. 20 (1999), pp. 511–530.
- [26] A. MIELKE AND P. SPRENGER, *Quasiconvexity at the boundary and a simple variational formulation of Agmon’s condition*, J. Elasticity, 51 (1998), pp. 23–41.
- [27] P. PEDREGAL, *Parametrized Measures and Variational Principles*, Birkhäuser, Basel, 1997.
- [28] F. RINDLER, *A local proof for the characterization of Young measures generated by sequences in BV* , J. Funct. Anal., 266 (2014), pp. 6335–6371.
- [29] T. ROUBÍČEK, *Relaxation in Optimization Theory and Variational Calculus*, W. de Gruyter, Berlin, 1997.
- [30] J. SOUČEK, *Spaces of functions on domain Ω , whose k -th derivatives are measures defined on $\bar{\Omega}$* , Časopis Pro Pěstování Matematiky, 97 (1972), pp. 10–46.
- [31] P. SPRENGER, *Quasikonvexität am Rande und Null-Lagrange-Funktionen in der nichtkonvexen Variationsrechnung*, Ph.D. thesis, Universität Hannover, 1996.
- [32] L. TARTAR, *Mathematical tools for studying oscillations and concentrations: From Young measures to H -measures and their variants*, in Multiscale Problems in Science and Technology: Challenges to Mathematical Analysis and Perspectives, N. Antoniċ et al., eds., Proceedings of the Conference on Multiscale Problems in Science and Technology, Dubrovnik, Croatia, 2000, Springer, Berlin, 2002.
- [33] M. VALADIER, *Young measures*, in Methods of Nonconvex Analysis, A. Cellina, ed., Lect. Notes Math. 1446, Springer, Berlin, 1990, pp. 152–188.
- [34] J. WARGA, *Optimal Control of Differential and Functional Equations*, Academic Press, New York, 1972.
- [35] L. C. YOUNG, *Generalized curves and the existence of an attained absolute minimum in the calculus of variations*, C. R. Soc. Sci. Lett. Varsovie, Classe III, 30 (1937), pp. 212–234.