

A nonlinear fractional partial integro-differential equation with nonlocal initial value conditions

Chenkuan Li¹  | Reza Saadati²  | Donal O'Regan³  | Radko Mesiar^{4,5}  |
Andrii Hrytsenko¹

¹Department of Mathematics and Computer Science, Brandon University, Brandon, Manitoba, Canada

²School of Mathematics, Iran University of Science and Technology, Tehran, Iran

³School of Mathematical and Statistical Science, University of Galway, Galway, Ireland

⁴Department of Mathematics, Faculty of Civil Engineering, Slovak University of Technology in Bratislava, Bratislava, Slovakia

⁵Institute of Information Theory and Automation, The Czech Academy of Sciences, Praha, Czech Republic

Correspondence

Reza Saadati, School of Mathematics, Iran University of Science and Technology, Narmak, Tehran 13114-16846, Iran.
Email: rsaadati@eml.cc

Chenkuan Li, Department of Mathematics and Computer Science, Brandon University, Brandon, Manitoba, Canada.
Email: lic@brandonu.ca

Communicated by: S. E. Mikhailov

Funding information

Natural Sciences and Engineering Research Council of Canada, Grant/Award Number: 2019-03907

In this work, we study a new nonlinear partial integro-differential equation with nonlocal initial value conditions and investigate the solutions of this equation. By considering an equivalent implicit integral equation via series, we prove the uniqueness of solutions of the equation by Babenko's approach, Banach's contraction principle, and the multivariable Mittag–Leffler function. We also demonstrate the application of our key theorem with an illustrative example.

KEYWORDS

Babenko's approach, Banach's contractive principle, multivariate Mittag–Leffler function, nonlinear partial integro-differential equation

MSC CLASSIFICATION

35A02, 35C15, 45E10, 26A33

1 | INTRODUCTION AND PRELIMINARIES

Let $b > 0$ and $c_0, c_1, c_2 \in C[0, b]$. We consider the uniqueness of solutions for the following NPDIE for $2 < \zeta \leq 3$, $0 < \varepsilon \leq 1$, and $\nu > 0$:

$$\begin{cases} \frac{\partial^\zeta \Phi}{\partial \theta^\zeta}(\theta, \sigma) + c_0(\sigma) \frac{\partial^\zeta \Phi}{\partial \theta^\zeta}(\theta, \sigma) + c_1(\sigma) \Phi(\theta, \sigma) + c_2(\sigma) I_\sigma^\nu \Phi(\theta, \sigma) = (\theta, \sigma, \Phi(\theta, \sigma)), \\ \Phi(0, \sigma) = \eta \int_0^1 \Phi(\theta, \sigma) d\theta, \quad \frac{\partial}{\partial \theta} \Phi(0, \sigma) = \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta, \quad \Phi_\theta''(0, \sigma) = 0, \end{cases} \quad (1.1)$$

where $(\theta, \sigma) \in [0, 1] \times [0, b]$, η is a constant, $\psi \in C[0, 1]$, and $\cdot : [0, 1] \times [0, b] \times R \rightarrow R$ satisfies certain conditions to be given later. Assume that $\nu \in R^+$ and $\sigma \in [0, b]$, the operator I_σ^ν is the partial Riemann–Liouville fractional integral of order

v w.r.t σ with initial point zero (see [1]),

$$(I_\sigma^v \Phi)(\theta, \sigma) = \frac{1}{\Gamma(v)} \int_0^\sigma (\sigma - \zeta)^{v-1} \Phi(\theta, \zeta) d\zeta,$$

and $\frac{\partial^\zeta}{\partial \theta^\zeta}$ is the partial Liouville–Caputo fractional derivative of order ζ with respect to $\theta \in [0, 1]$

$$\left(\frac{\partial^\zeta}{\partial \theta^\zeta} \Phi \right) (\theta, \sigma) = \frac{1}{\Gamma(3 - \zeta)} \int_0^\theta (\theta - s)^{2-\zeta} \Phi_s^{(3)}(s, \sigma) ds, \quad 2 < \zeta \leq 3,$$

and similarly,

$$\left(\frac{\partial^\varepsilon}{\partial \theta^\varepsilon} \Phi \right) (\theta, \sigma) = \frac{1}{\Gamma(1 - \varepsilon)} \int_0^\theta (\theta - s)^{-\varepsilon} \Phi'_s(s, \sigma) ds, \quad 0 < \varepsilon \leq 1.$$

A function Φ is said to be a solution of Equation (1.1) if it satisfies the equation and its three initial conditions.

If $\psi(\theta) = \eta$ for all $\theta \in [0, 1]$, then Equation (1.1) is

$$\begin{cases} \frac{\partial^\zeta}{\partial \theta^\zeta} \Phi(\theta, \sigma) + c_0(\sigma) \frac{\partial^\varepsilon}{\partial \theta^\varepsilon} \Phi(\theta, \sigma) + c_1(\sigma) \Phi(\theta, \sigma) + c_2(\sigma) I_\sigma^v \Phi(\theta, \sigma) = F(\theta, \sigma, \Phi(\theta, \sigma)), \\ \Phi(0, \sigma) = \frac{\partial}{\partial \theta} \Phi(0, \sigma) = \eta \int_0^1 \Phi(\theta, \sigma) d\theta, \quad \Phi_t''(0, \sigma) = 0. \end{cases}$$

It follows from [2] that

$$(I_\sigma^0 \Phi)(\theta, \sigma) = \Phi(\theta, \sigma).$$

It is well known that fractional partial differential equations have been used in many fields such as quantum mechanics, fluid dynamics, and plasma physics [3]. In 2016, Petitta [4] considered the following boundary value problem and obtained solutions for this equation and investigated the uniqueness and regularity of the solutions:

$$\begin{cases} \mathcal{L}^s \Phi = \mu \text{ in } \Omega, \quad 0 < s < 1, \\ \Phi(\sigma) = 0, \text{ on } R^n \setminus \Omega, \end{cases}$$

where Ω is a bounded domain of R^n , μ is a bounded Radon measure on Ω , and \mathcal{L}^s is given as

$$\mathcal{L}^s \Phi(\sigma) = PV \int_{R^n} (\Phi(\sigma) - \Phi(\sigma + y)) K(y) dy,$$

and K is a non-negative kernel satisfying

$$\frac{\lambda}{|y|^{n+2s}} \leq K(y) \leq \frac{\Lambda}{|y|^{n+2s}}, \quad 0 < \lambda \leq \Lambda.$$

This class of kernels contains the fractional Laplacian $(-\Delta)^s$ as a particular case, that is,

$$(-\Delta)^s \Phi(\sigma) := c_{n,s} PV \int_{R^n} \frac{\Phi(\sigma) - \Phi(\sigma + y)}{|y|^{n+2s}} dy,$$

where

$$c_{n,s} = \frac{4^s s(s-1) \Gamma\left(\frac{n+2s}{2}\right)}{\pi^{n/2} \Gamma(2-s)}.$$

In [5], the author extended the fractional Laplacian $(-\Delta)^s \Phi$ over a space to all values of $s > 0$ and $s \neq 1, 2, \dots$, based on the normalization in distribution theory (in the sense of Schwartz), Pizzetti's formula, and surface integrals in R^n . Recently, Mahor et al. [6] investigated several fractional equations and obtained analytical solutions using the fractional

Fourier transform and MLF for these equations. The fractional heat-diffusion equation is one of these equations, which is considered as follows:

$$\begin{cases} \frac{\partial^\zeta}{\partial \theta^\zeta} \phi(\sigma, \theta) = h \frac{\partial^{2\zeta}}{\partial \theta^{2\zeta}} \phi(\sigma, \theta), \sigma \in R, \theta \in R^+, 0 < \zeta < 1, \\ \phi(\sigma, 0) = \psi(\sigma), \end{cases}$$

where $\phi(\sigma, \theta)$ is the temperature function and $h \in R^+$ is the thermal conductivity.

In the following, we introduce the multivariable Mittag–Leffler function (MMLF) and Babenko's approach (B-A); we solve the fractional kinetic equation in detail. In this paper, we consider the Banach space $C([0, 1] \times [0, b])$ with the norm

$$\|\Phi\| = \sup_{\theta \in [0, 1], \sigma \in [0, b]} |\Phi(\theta, \sigma)| \text{ for } \Phi \in C([0, 1] \times [0, b]).$$

Definition 1 ([7]). The two-parameter Mittag–Leffler (M-L) function is represented by the following series:

$$E_{\zeta, \varepsilon}(z) = \sum_{h=0}^{\infty} \frac{z^h}{\Gamma(\zeta h + \varepsilon)},$$

where $\zeta, \varepsilon > 0$ and $z \in C$ (the complex plane). The MMLF is defined as follows:

$$E_{(\zeta_1, \dots, \zeta_m), \varepsilon}(z_1, \dots, z_m) = \sum_{h=0}^{\infty} \sum_{\substack{h_1 + \dots + h_m = h \\ h_1 \geq 0, \dots, h_m \geq 0}} \binom{h}{h_1, \dots, h_m} \frac{z_1^{h_1} \dots z_m^{h_m}}{\Gamma(\zeta_1 h_1 + \dots + \zeta_m h_m + \varepsilon)},$$

where $\zeta_i, \varepsilon > 0$, for $i = 1, 2, \dots, m$ and

$$\binom{h}{h_1, \dots, h_m} = \frac{h!}{h_1! \dots h_m!}.$$

One of the useful tools for dealing with differential equations with initial conditions as well as integral equations is B-A. This method is generally the same as the Laplace transform, while working on equations with constant coefficients. However, this method is widely used for differential and integral equations with continuous variable coefficients and boundary value problems [8]. In the following, to show some details of this approach, we will deduce the solution to the following fractional kinetic equation investigated by Saxena and Kalla in [9]

$$N(\theta) - N_0 \psi(\theta) = -c^\nu I_t^\nu N(\theta), \nu > 0, \quad (1.2)$$

where $N(\theta)$ denotes the number density of a given species at time θ , $N_0 = N(0)$ is the number density of that species at time $\theta = 0$ and c is a positive constant.

Clearly, Equation (1.2) can be written as

$$(1 + c^\nu I_\theta^\nu) N(\theta) = N_0 \psi(\theta).$$

Treating the factor $(1 + c^\nu I_\theta^\nu)$ as a variable, we derive from B-A

$$\begin{aligned} N(\theta) &= N_0 (1 + c^\nu I_\theta^\nu)^{-1} \psi(\theta) = N_0 \sum_{h=0}^{\infty} (-1)^h (c^\nu I_\theta^\nu)^h \psi(\theta) \\ &= N_0 \sum_{h=0}^{\infty} (-1)^h c^{h\nu} I_\theta^{h\nu} \psi(\theta) \\ &= N_0 \psi(\theta) + N_0 \sum_{h=1}^{\infty} (-1)^h c^{h\nu} \frac{1}{\Gamma(h\nu)} \int_0^\theta (\theta - \zeta)^{h\nu-1} \psi(\zeta) d\zeta \end{aligned}$$

$$\begin{aligned}
 &= N_0\psi(\theta) + N_0 \sum_{h=0}^{\infty} (-1)^{h+1} c^{hv+v} \frac{1}{\Gamma(hv+v)} \int_0^\theta (\theta-\zeta)^{hv+v-1} \psi(\zeta) d\zeta \\
 &= N_0\psi(\theta) - N_0 c^v \int_0^\theta (\theta-\zeta)^{v-1} \sum_{h=0}^{\infty} (-1)^h c^{hv} \frac{(\theta-\zeta)^{hv}}{\Gamma(hv+v)} \psi(\zeta) d\zeta \\
 &= N_0\psi(\theta) - N_0 c^v \int_0^\theta (\theta-\zeta)^{v-1} E_{v,v}(-c^v(\theta-\zeta)^v) \psi(\zeta) d\zeta,
 \end{aligned}$$

which is well defined. We should note that this is indeed Abel's equation of the second kind. There are some recent studies on several general classes of fractional-order kinetic equations using various approaches [6, 10–12]. In addition, researchers have investigated the existence and uniqueness for various types of fractional integro-differential equations, with boundary conditions and the Hilfer derivative, via different methods [13–16].

The main goal of this paper is to transform Equation (1.1) into an equivalent implicit integral equation in the series using B-A, which is derived in Section 2. Also, with the help of Banach's contraction principle (BCP), the uniqueness of the solutions in $C([0, 1] \times [0, b])$ is studied. In Section 3, we present the application of proven results with an example, and finally, we summarize the entire work in Section 4.

2 | MAIN RESULTS

Theorem 2. *Let $b > 0$, $2 < \zeta \leq 3$, $0 < \varepsilon \leq 1$, $v > 0$, η be a constant and $c_0, c_1, c_2, \psi \in C[0, b]$. We assume $F : [0, 1] \times [0, b] \times R \rightarrow R$ is a bounded and continuous function, and*

$$2(|\eta| + m_0|\eta| + M) E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2 b^v) < 1,$$

where

$$m_i = \max_{\sigma \in [0, b]} |c_i(\sigma)|, \text{ and } M = \max_{\sigma \in [0, b]} |\psi(\sigma)|,$$

for $i = 0, 1, 2$. Then Φ is a solution of Equation (1.1) if and only if it satisfies the following implicit integral equation in the space $C([0, 1] \times [0, b])$:

$$\begin{aligned}
 \Phi(\theta, \sigma) &= \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\cdot I_t^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+\zeta} (c_2(\sigma) I_\sigma^v)^{h_3} F(\theta, \sigma, \Phi(\theta, \sigma)) \\
 &+ \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\cdot \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3}}{\Gamma((\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+1)} \eta \int_0^1 (c_2(\sigma) I_\sigma^v)^{h_3} \Phi(\theta, \sigma) d\theta \\
 &+ \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\cdot \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+\zeta-\varepsilon}}{\Gamma((\zeta-\beta)h_1+\zeta h_2+\zeta h_3+\zeta-\beta+1)} \eta \int_0^1 (c_2(\sigma) I_\sigma^v)^{h_3} c_0(\sigma) \Phi(\theta, \sigma) d\theta \\
 &+ \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\cdot \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+1}}{\Gamma((\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+2)} \int_0^1 \psi(\theta) (c_2(\sigma) I_\sigma^v)^{h_3} \Phi(\theta, \sigma) d\theta.
 \end{aligned} \tag{2.1}$$

Proof. It can be clearly seen that

$$\begin{aligned} I_{\theta}^{\zeta} \frac{c \partial^{\zeta}}{\partial \theta^{\zeta}} \Phi(\theta, \sigma) &= \Phi(\theta, \sigma) - \Phi(0, \sigma) - \Phi'_{\theta}(0, \sigma) \theta - \frac{\Phi''_{\theta}(0, \sigma)}{2!} \theta^2 \\ &= \Phi(\theta, \sigma) - \eta \int_0^1 \Phi(\theta, \sigma) d\theta - \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta, \end{aligned}$$

and since $0 < \varepsilon \leq 1$

$$\begin{aligned} I_{\theta}^{\zeta} c_0(\sigma) \frac{c \partial^{\varepsilon}}{\partial \theta^{\varepsilon}} \Phi(\theta, \sigma) &= c_0(\sigma) I_{\theta}^{\zeta-\beta} I_{\theta}^{\zeta} \frac{c \partial^{\varepsilon}}{\partial \theta^{\varepsilon}} \Phi(\theta, \sigma) \\ &= c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} (\Phi(\theta, \sigma) - \Phi(0, \sigma)) \\ &= c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} \Phi(\theta, \sigma) - \frac{c_0(\sigma) \theta^{\zeta-\beta} \eta}{\Gamma(\zeta - \beta + 1)} \int_0^1 \Phi(\theta, \sigma) d\theta. \end{aligned}$$

Applying the operator I_{θ}^{ζ} to the equation

$$\frac{c \partial^{\zeta}}{\partial \theta^{\zeta}} \Phi(\theta, \sigma) + c_0(\sigma) \frac{c \partial^{\varepsilon}}{\partial \theta^{\varepsilon}} \Phi(\theta, \sigma) + c_1(\sigma) \Phi(\theta, \sigma) + c_2(\sigma) I_{\theta}^{\zeta} \Phi(\theta, \sigma) = F(\theta, \sigma, \Phi(\theta, \sigma)),$$

we get

$$\begin{aligned} &(1 + c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} + c_1(\sigma) I_{\theta}^{\zeta} + c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta}) \Phi(\theta, \sigma) \\ &= I_{\theta}^{\zeta} F(\theta, \sigma, \Phi(\theta, \sigma)) + \left(1 + \frac{c_0(\sigma) \theta^{\zeta-\varepsilon}}{\Gamma(\zeta - \varepsilon + 1)}\right) \eta \int_0^1 \Phi(\theta, \sigma) d\theta + \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta. \end{aligned}$$

Using B-A, we obtain

$$\begin{aligned} \Phi(\theta, \sigma) &= (1 + c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} + c_1(\sigma) I_{\theta}^{\zeta} + c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta})^{-1} \\ &\quad \cdot \left[I_{\theta}^{\zeta} F(\theta, \sigma, \Phi(\theta, \sigma)) + \left(1 + \frac{c_0(\sigma) \theta^{\zeta-\varepsilon}}{\Gamma(\zeta - \varepsilon + 1)}\right) \eta \int_0^1 \Phi(\theta, \sigma) d\theta + \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta \right] \\ &= \sum_{h=0}^{\infty} (-1)^h (c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} + c_1(\sigma) I_{\theta}^{\zeta} + c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta})^h \\ &\quad \cdot \left[I_{\theta}^{\zeta} F(\theta, \sigma, \Phi(\theta, \sigma)) + \left(1 + \frac{c_0(\sigma) \theta^{\zeta-\varepsilon}}{\Gamma(\zeta - \varepsilon + 1)}\right) \eta \int_0^1 \Phi(\theta, \sigma) d\theta + \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta \right] \\ &= \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} (c_0(\sigma) I_{\theta}^{\zeta-\varepsilon})^{h_1} (c_1(\sigma) I_{\theta}^{\zeta})^{h_2} (c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta})^{h_3} \\ &\quad \cdot \left[I_{\theta}^{\zeta} F(\theta, \sigma, \Phi(\theta, \sigma)) + \left(1 + \frac{c_0(\sigma) \theta^{\zeta-\varepsilon}}{\Gamma(\zeta - \varepsilon + 1)}\right) \eta \int_0^1 \Phi(\theta, \sigma) d\theta + \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta \right]. \end{aligned}$$

Clearly,

$$(c_0(\sigma) I_{\theta}^{\zeta-\varepsilon})^{h_1} = c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} \cdots c_0(\sigma) I_{\theta}^{\zeta-\varepsilon} = c_0^{h_1}(\sigma) I_{\theta}^{(\zeta-\varepsilon)h_1}$$

and $(c_1(\sigma) I_{\theta}^{\zeta})^{h_2} = c_1^{h_2}(\sigma) I_{\theta}^{\zeta h_2}$. As for the term $(c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta})^{h_3}$, we have

$$(c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta})^{h_3} = (c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta}) (c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta}) \cdots (c_2(\sigma) I_{\theta}^{\zeta} I_{\theta}^{\zeta}) = I_{\theta}^{\zeta h_3} (c_2(\sigma) I_{\theta}^{\zeta})^{h_3}.$$

Therefore,

$$\begin{aligned} \Phi(\theta, \sigma) &= \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) I_{\theta}^{(\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3} (c_2(\sigma) I_{\theta}^{\zeta})^{h_3} \\ &\quad \cdot \left[I_{\theta}^{\zeta} F(\theta, \sigma, \Phi(\theta, \sigma)) + \left(1 + \frac{c_0(\sigma) \theta^{\zeta-\varepsilon}}{\Gamma(\zeta - \varepsilon + 1)}\right) \eta \int_0^1 \Phi(\theta, \sigma) d\theta + \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta \right] \end{aligned}$$

$$\begin{aligned}
 &= \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot I_{\theta}^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+\zeta} (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} F(\theta, \sigma, \Phi(\theta, \sigma)) \\
 &\quad + \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3}}{\Gamma((\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+1)} \eta \int_0^1 (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} \Phi(\theta, \sigma) d\theta \\
 &\quad + \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot \frac{I_{\theta}^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3} \theta^{\zeta-\varepsilon}}{\Gamma(\zeta-\beta+1)} \eta \int_0^1 (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} c_0(\sigma) \Phi(\theta, \sigma) d\theta \\
 &\quad + \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot I_{\theta}^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3} \theta \int_0^1 \psi(\theta) (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} \Phi(\theta, \sigma) d\theta.
 \end{aligned}$$

Using

$$\begin{aligned}
 I_{\theta}^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3} \theta^{\zeta-\varepsilon} &= \frac{\Gamma(\zeta-\varepsilon+1)}{\Gamma((\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+\zeta-\beta+1)} \theta^{(\zeta-\beta)h_1+\zeta h_2+\zeta h_3+\zeta-\beta}, \\
 I_{\theta}^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3} \theta &= \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+1}}{\Gamma((\zeta-\beta)h_1+\zeta h_2+\zeta h_3+2)},
 \end{aligned}$$

we have

$$\begin{aligned}
 \Phi(\theta, \sigma) &= \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot I_{\theta}^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+\zeta} (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} F(\theta, \sigma, \Phi(\theta, \sigma)) \\
 &\quad + \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3}}{\Gamma((\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+1)} \eta \int_0^1 (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} \Phi(\theta, \sigma) d\theta \\
 &\quad + \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+\zeta-\varepsilon}}{\Gamma((\zeta-\beta)h_1+\zeta h_2+\zeta h_3+\zeta-\beta+1)} \eta \int_0^1 (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} c_0(\sigma) \Phi(\theta, \sigma) d\theta \\
 &\quad + \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\
 &\quad \cdot \frac{\theta^{(\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+1}}{\Gamma((\zeta-\varepsilon)h_1+\zeta h_2+\zeta h_3+2)} \int_0^1 \psi(\theta) (c_2(\sigma) I_{\sigma}^{\nu})^{h_3} \Phi(\theta, \sigma) d\theta.
 \end{aligned}$$

The function $\Phi(\theta, \sigma)$ satisfies the initial conditions: First $\Phi(0, \sigma) = \eta \int_0^1 \Phi(\theta, \sigma) d\sigma$, which is the term

$$(-1)^0 \sum_{\substack{h_1+h_2+h_3=0 \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{0}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \cdot \frac{\theta^{(\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3}}{\Gamma((\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + 1)} \eta \int_0^1 (c_2(\sigma) I_\sigma^\nu)^{h_3} \Phi(\theta, \sigma) d\theta,$$

and the rest being zero by setting $\theta = 0$. Similarly, $\frac{\partial}{\partial \theta} \Phi(0, \sigma) = \int_0^1 \psi(\theta) \Phi(\theta, \sigma) d\theta$, $\Phi'_\theta(0, \sigma) = 0$, by noting that $\zeta - \varepsilon > 1$. Hence Φ is a solution of Equation (1.1) if and only if it satisfies Equation (2.1). Next, in $C([0, 1] \times [0, b])$, we show the convergence of the right side of Equation (2.1), which is a series. Since $c_0, c_1, c_2 \in C[0, b]$, we let $m_i = \max_{\sigma \in [0, b]} |c_i(\sigma)|$, for $i = 0, 1, 2$. Then,

$$\begin{aligned} \|\Phi\| &\leq \sum_{h=0}^{\infty} \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} m_0^{h_1} m_1^{h_2} m_2^{h_3} \\ &\cdot \frac{b^{\nu h_3}}{\Gamma((\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + \zeta + 1) \Gamma(\nu h_3 + 1)} \sup_{(\theta, \sigma) \in [0, 1] \times [0, b]} |F(\theta, \sigma, \Phi(\theta, \sigma))| \\ &+ \sum_{h=0}^{\infty} \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} m_0^{h_1} m_1^{h_2} m_2^{h_3} \\ &\cdot \frac{b^{\nu h_3} |\eta|}{\Gamma((\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + 1) \Gamma(\nu h_3 + 1)} \|\Phi\| \\ &+ \sum_{h=0}^{\infty} \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} m_0^{h_1+1} m_1^{h_2} m_2^{h_3} \\ &\cdot \frac{b^{\nu h_3} |\eta|}{\Gamma((\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + \zeta - \varepsilon + 1) \Gamma(\nu h_3 + 1)} \|\Phi\| \\ &+ \sum_{h=0}^{\infty} \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} m_0^{h_1} m_1^{h_2} m_2^{h_3} \\ &\cdot \frac{M b^{\nu h_3}}{\Gamma((\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + 2) \Gamma(\nu h_3 + 1)} \|\Phi\|, \end{aligned}$$

where $M = \max_{\sigma \in [0, b]} |\psi(\sigma)|$. Applying the property $\Gamma(\nu h_3 + 1) \geq 1/2$, we deduce

$$\begin{aligned} \|\Phi\| &\leq 2E_{(\zeta-\varepsilon, \zeta, \zeta), \zeta+1} (m_0, m_1, m_2 b^\nu) \sup_{(\theta, \sigma) \in [0, 1] \times [0, b]} |F(\theta, \sigma, \Phi(\theta, \sigma))| \\ &\quad + 2|\eta| E_{(\zeta-\varepsilon, \zeta, \zeta), 1} (m_0, m_1, m_2 b^\nu) \|\Phi\| \\ &\quad + 2m_0 |\eta| E_{(\zeta-\varepsilon, \zeta, \zeta), \zeta-\varepsilon+1} (m_0, m_1, m_2 b^\nu) \|\Phi\| \\ &\quad + 2M E_{(\zeta-\varepsilon, \zeta, \zeta), 2} (m_0, m_1, m_2 b^\nu) \|\Phi\| \\ &\leq 2E_{(\zeta-\varepsilon, \zeta, \zeta), \zeta+1} (m_0, m_1, m_2 b^\nu) \sup_{(\theta, \sigma) \in [0, 1] \times [0, b]} |F(\theta, \sigma, \Phi(\theta, \sigma))| \\ &\quad + 2(|\eta| + m_0 |\eta| + M) E_{(\zeta-\varepsilon, \zeta, \zeta), 1} (m_0, m_1, m_2 b^\nu) \|\Phi\|. \end{aligned}$$

From $q = 1 - 2(|\eta| + m_0 |\eta| + M) E_{(\zeta-\varepsilon, \zeta, \zeta), 1} (m_0, m_1, m_2 b^\nu) > 0$, we have

$$\|\Phi\| \leq \frac{2}{q} E_{(\zeta-\varepsilon, \zeta, \zeta), \zeta+1} (m_0, m_1, m_2 b^\nu) \sup_{(\theta, \sigma) \in [0, 1] \times [0, b]} |F(\theta, \sigma, \Phi(\theta, \sigma))| < +\infty,$$

since F is a bounded function. This implies that the series is well defined in the space $C([0, 1] \times [0, b])$. This completes the proof. □

In the following, we present and prove the uniqueness of the solutions of Equation (1.1) in the form of a theorem.

Theorem 3. Let $b > 0, 2 < \zeta \leq 3, 0 < \varepsilon \leq 1, \nu > 0, \eta$ be a constant and $c_0, c_1, c_2, f \in C[0, b]$. We assume the continuous and bounded function $F : [0, 1] \times [0, b] \times R \rightarrow R$ satisfies the following Lipschitz condition with a constant $L \geq 0$:

$$|F(\theta, \sigma, y_1) - F(\theta, \sigma, y_2)| \leq L|y_1 - y_2|, (\theta, \sigma) \in [0, 1] \times [0, b], y_1, y_2 \in R.$$

In addition, we suppose

$$2(L + |\eta| + m_0|\eta| + M)E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2b^\nu) < 1,$$

where

$$m_i = \max_{\sigma \in [0, b]} |c_i(\sigma)|, \text{ and } M = \max_{\sigma \in [0, b]} |\psi(\sigma)|,$$

for $i = 0, 1, 2$. Then Equation (1.1) has a unique solution in the space $C([0, 1] \times [0, b])$.

Proof. We consider the nonlinear mapping T over the space $C([0, 1] \times [0, b])$ and define it as follows:

$$\begin{aligned} (T\Phi)(\theta, \sigma) &= \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\ &\cdot I_t^{(\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + \zeta} (c_2(\sigma) I_\sigma^\nu)^{h_3} F(\theta, \sigma, \Phi(\theta, \sigma)) \\ &+ \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\ &\cdot \frac{\theta^{(\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3}}{\Gamma((\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + 1)} \eta \int_0^1 (c_2(\sigma) I_\sigma^\nu)^{h_3} \Phi(\theta, \sigma) d\theta \\ &+ \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\ &\cdot \frac{\theta^{(\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + \zeta - \varepsilon}}{\Gamma((\zeta-\beta)h_1 + \zeta h_2 + \zeta h_3 + \zeta - \beta + 1)} \eta \int_0^1 (c_2(\sigma) I_\sigma^\nu)^{h_3} c_0(\sigma) \Phi(\theta, \sigma) d\theta \\ &+ \sum_{h=0}^{\infty} (-1)^h \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} c_0^{h_1}(\sigma) c_1^{h_2}(\sigma) \\ &\cdot \frac{\theta^{(\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + 1}}{\Gamma((\zeta-\varepsilon)h_1 + \zeta h_2 + \zeta h_3 + 2)} \int_0^1 \psi(\theta) (c_2(\sigma) I_\sigma^\nu)^{h_3} \Phi(\theta, \sigma) d\theta. \end{aligned}$$

It follows from the proof of Theorem 2 that $T\Phi \in C([0, 1] \times [0, b])$. We now show that T is contractive. Indeed, similar to that in the proof of Theorem 2,

$$\begin{aligned} \|T\Phi - T\vartheta\| &\leq 2E_{(\zeta-\varepsilon, \zeta, \zeta), \zeta+1}(m_0, m_1, m_2b^\nu) \sup_{(\theta, \sigma) \in [0, 1] \times [0, b]} |F(\theta, \sigma, \Phi(\theta, \sigma)) - F(\theta, \sigma, \vartheta(\theta, \sigma))| \\ &+ 2|\eta|E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2b^\nu) \|\Phi - \vartheta\| \\ &+ 2m_0|\eta|E_{(\zeta-\varepsilon, \zeta, \zeta), \zeta-\varepsilon+1}(m_0, m_1, m_2b^\nu) \|\Phi - \vartheta\| \\ &+ 2ME_{(\zeta-\varepsilon, \zeta, \zeta), 2}(m_0, m_1, m_2b^\nu) \|\Phi - \vartheta\| \\ &\leq 2LE_{(\zeta-\varepsilon, \zeta, \zeta), \zeta+1}(m_0, m_1, m_2b^\nu) \|\Phi - \vartheta\| \\ &+ 2(|\eta| + m_0|\eta| + M)E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2b^\nu) \|\Phi - \vartheta\| \\ &\leq 2(L + |\eta| + m_0|\eta| + M)E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2b^\nu) \|\Phi - \vartheta\|. \end{aligned}$$

Since $2(L + |\eta| + m_0|\eta| + M)E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2b^\nu) < 1$, Equation (1.1) has a unique solution in the space $C([0, 1] \times [0, b])$ by BCP. □

3 | EXAMPLE

Example 4. We consider an NPDIE-NIVC as follows:

$$\begin{cases} \frac{\partial^{2.5}}{\partial \theta^{2.5}} \Phi(\theta, \sigma) + \sin \sigma \frac{\partial^{0.5}}{\partial \theta^{0.5}} \Phi(\theta, \sigma) + \frac{\sigma^2}{1+\sigma^2} \Phi(\theta, \sigma) + \cos \sigma I_\sigma \Phi(\theta, \sigma) = \frac{1}{21} \sin |\Phi(\theta, \sigma)|, \\ \Phi(0, \sigma) = \frac{1}{40} \int_0^1 \Phi(\theta, \sigma) d\theta, \quad \frac{\partial}{\partial \theta} \Phi(0, \sigma) = \frac{1}{22} \int_0^1 \theta \Phi(\theta, \sigma) d\theta, \quad \Phi'_\theta(0, \sigma) = 0. \end{cases}$$

Then, it has a unique solution in the space $C([0, 1] \times [0, 1])$.

Proof. Clearly, $m_0 = m_1 = m_2 = 1$, and $F(\theta, \sigma, y) = \frac{1}{21} \sin |y|$, is a continuous and bounded function satisfying

$$|F(\theta, \sigma, y_1) - F(\theta, \sigma, y_2)| \leq \frac{1}{21} |y_1 - y_2|.$$

Thus, $L = 1/21$. Moreover, $M = 1/22$ and $\eta = 1/40$. Finally, we note

$$\begin{aligned} & 2(L + |\eta| + m_0|\eta| + M) E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2 b^v) \\ &= 2(1/21 + 1/40 + 1/40 + 1/22) E_{(2, 2.5, 2.5), 1}(1, 1, 1) \\ &\leq \frac{3}{10} E_{(2, 2.5, 2.5), 1}(1, 1, 1), \end{aligned}$$

and

$$\begin{aligned} E_{(2, 2.5, 2.5), 1}(1, 1, 1) &= \sum_{h=0}^{\infty} \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} \frac{1}{\Gamma(2h_1 + 2.5h_2 + 2.5h_3 + 1)} \\ &\leq \sum_{h=0}^{\infty} \frac{3^h}{(2h)!} = \cosh(\sqrt{3}) \approx 2.9146, \end{aligned}$$

since

$$\begin{aligned} \sum_{\substack{h_1+h_2+h_3=h \\ h_1 \geq 0, h_2 \geq 0, h_3 \geq 0}} \binom{h}{h_1, h_2, h_3} &= 3^h, \\ \frac{1}{\Gamma(2h_1 + 2.5h_2 + 2.5h_3 + 1)} &\leq \frac{1}{\Gamma(2h + 1)} = \frac{1}{(2h)!}. \end{aligned}$$

Hence,

$$2(L + |\eta| + m_0|\eta| + M) E_{(\zeta-\varepsilon, \zeta, \zeta), 1}(m_0, m_1, m_2 b^v) < 1.$$

By Theorem 3, the NPDIE-NIVC has a unique solution in the space $C([0, 1] \times [0, 1])$. \square

4 | CONCLUSION

In this article, we have considered the NPDIE with nonlocal initial value conditions. Our main work was to investigate the uniqueness of solutions for this equation using an implicit integral equation, B-A, MMLF, and BCP. In the final part, we have presented the numerical results of our key theorem by giving an illustrative example.

AUTHOR CONTRIBUTIONS

All authors conceived of the study, participated in its design and coordination, drafted the manuscript, participated in the sequence alignment, and read and approved the final manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no competing interests.

ORCID

Chenkuan Li  <https://orcid.org/0000-0001-7098-8059>

Reza Saadati  <https://orcid.org/0000-0002-6770-6951>

Donal O'Regan  <https://orcid.org/0000-0002-4096-1469>

Radko Mesiar  <https://orcid.org/0000-0001-6503-949X>

REFERENCES

1. A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and applications of fractional differential equations*, Elsevier, North-Holland, 2006.
2. C. Li, *Several results of fractional derivatives in $D'(R_+)$* , *Fract. Calc. Appl. Anal.* **18** (2015), 192–207.
3. S. Momani and Z. Odibat, *Analytical approach to linear fractional partial differential equations arising in fluid mechanics*, *Phys. Lett. A* **355** (2006), 271–279.
4. F. Petitta, *Some remarks on the duality method for integro-differential equations with measure data*, *Adv. Nonlinear Stud.* **16** (2016), 115–124.
5. C. Li, *On the generalized fractional Laplacian*, *Fract. Calc. Appl. Anal.* **24** (2021), 1797–1830, DOI 10.1515/fca-2021-0078.
6. T. C. Mahor, R. Mishra, and R. Jain, *Analytical solutions of linear fractional partial differential equations using fractional Fourier transform*, *J. Comput. Appl. Math.* **385** (2021), 113202.
7. S. B. Hadid and Y. F. Luchko, *An operational method for solving fractional differential equations of an arbitrary real order*, *Panamer. Math. J.* **6** (1996), 57–73.
8. Y. I. Babenko, *Heat and mass transfer*, Khimiya, Leningrad, 1986. (in Russian).
9. R. K. Saxena and S. L. Kalla, *On the solution of certain fractional kinetic equations*, *Appl. Math. Comput.* **199** (2008), 504–511, DOI 10.1016/j.amc.2007.10.005.
10. T. T. Phong and L. D. Long, *Well-posed results for nonlocal fractional parabolic equation involving Caputo-Fabrizio operator*, *J. Math. Computer Sci.* **26** (2022), no. 4, 357–367, DOI 10.22436/jmcs.026.04.04.
11. P. O. Mohammed, H. M. Srivastava, J. L. G. Guirao, et al., *Existence of solutions for a class of nonlinear fractional difference equations of the Riemann-Liouville type*, *Adv. Cont. Discr. Mod.* **2022** (2022), 32, DOI 10.1186/s13662-022-03705-9.
12. H. M. Srivastava, B. J. González, and E. R. Negrín, *An operational calculus for a Mehler-Fock type index transform on distributions of compact support*, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.* **117** (2023), 3, DOI 10.1007/s13398-022-01335-0.
13. M. I. Youssef, *Generalized fractional delay functional equations with Riemann-Stieltjes and infinite point nonlocal conditions*, *J. Math. Computer Sci.* **24** (2022), no. 1, 33–48, DOI 10.22436/jmcs.024.01.04.
14. T. Abdeljawad, P. O. Mohammed, H. M. Srivastava, E. Al-Sarairah, A. Kashuri, and K. Nonlaopon, *Some novel existence and uniqueness results for the Hilfer fractional integro-differential equations with non-instantaneous impulsive multi-point boundary conditions and their application*, *AIMS Math.* **8** (2022), 3469–3483.
15. S. Kumar, R. H. Pandey, H. M. Srivastava, and G. N. Singh, *A convergent collocation approach for generalized fractional integro-differential equations using Jacobi poly-fractionals*, *Mathematics* **9** (2021), 979, 1–17.
16. M. R. Ali, A. R. Hadhoud, and H. M. Srivastava, *Solution of fractional Volterra-Fredholm integro-differential equations under mixed boundary conditions by using the HOBW method*, *Adv. Differ. Equ.* **2019** (2019), 115, 1–14.

How to cite this article: C. Li, R. Saadati, D. O'Regan, R. Mesiar, and A. Hrytsenko, *A nonlinear fractional partial integro-differential equation with nonlocal initial value conditions*, *Math. Meth. Appl. Sci.* **46** (2023), 17010–17019, DOI 10.1002/mma.9486.